


DEPARTMENT OF BASIC SCIENCE
ENGINEERING MATHEMATICS-III



Prepared by:-SUMITRA SAHU

RAJAKISHORE CHANDRA ACADEMY OF TECHNOLOGY
NILGIRI, Balasore (Odisha)

CONTENT

CHAPTER-1	MATRICES	2-7
CHAPTER-2	COMPLEXNUMBER	8-10
CHAPTER-2	LINEARDIFFERENTIALEQUATION	11-24
CHAPTER-3	LAPLACETRANSFORMATION	25-42
CHAPTER-4	FOURIERSERIES	43-52
CHAPTER-5	FINITEDIFFERENCE&INTERPOLATION	53-64
CHAPTER-6	NUMERICALSOLUTIONOFEQUATION	65-71

CHAPTER–1

MATRICES

Minor – Minor is the determinate value which is obtained by deleting row & column of the particular element and denoted by the symbol M_{ij} , i-rows j-column.

$$\text{Ex: } \begin{vmatrix} 2 & 1 & 3 \\ 4 & 2 & 8 \\ 5 & 6 & 1 \end{vmatrix}$$

$$M_{21} = \begin{vmatrix} 1 & 3 \\ 6 & 1 \end{vmatrix} = 1 - 18 = -17$$

$$M_{32} = \begin{vmatrix} 2 & 3 \\ 4 & 8 \end{vmatrix} = 16 - 12 = 4$$

Upper triangular Matrix – A matrix is said to be upper triangular if the elements below the main diagonal are zeros.

$$\text{Ex: } \begin{vmatrix} 1 & 5 \\ 0 & 3 \\ 0 & 8 \end{vmatrix}$$

Elementary transformations: – The following operations three of which refer to rows are known as elementary transformations.

- I. The interchange of any two rows ($R_i \leftrightarrow R_j$)
- II. The multiplication of any row by a non-zero scalar (kR_i)
- III. The addition of a constant multiple of the elements of any row to the corresponding elements of any other row ($R_i + kR_j$)

Equivalent matrix – Two matrices A and B are said to be equivalent if one can be obtained from the other by a sequence of elementary transformations.

Rank of a matrix: A matrix is said to be of rank 'r' if

- (i) It has at least one non-zero minor of order 'r'
- (ii) Every minor of order higher than 'r' vanishes.

The rank of a matrix A shall be denoted by the symbol $\rho(A)$.

Working Rule :**Step–I:** Convert the matrix to the upper triangular form.**Step–II:** Then the no. of non-zero rows is the rank of the matrix**Example–1:**

Find the rank of the matrix

$$A = \begin{bmatrix} 3 & -1 & 2 \\ -6 & 2 & 4 \\ -3 & 1 & 2 \end{bmatrix}$$

Solution:

$$A = \begin{bmatrix} 3 & -1 & 2 \\ -6 & 2 & 4 \\ -3 & 1 & 2 \end{bmatrix}$$

$$\sim \begin{bmatrix} 3 & -1 & 2 \\ 0 & 0 & 8 \\ -3 & 1 & 2 \end{bmatrix} \quad \begin{matrix} R_2 \rightarrow R_2 + 2R_1 \\ R_3 \rightarrow R_3 + R_1 \end{matrix}$$

$$\sim \begin{bmatrix} 3 & -1 & 2 \\ 0 & 0 & 8 \\ 0 & 0 & 4 \end{bmatrix}$$

$$\sim \begin{bmatrix} 3 & -1 & 2 \\ 0 & 0 & 8 \\ 0 & 0 & 0 \end{bmatrix} \quad \begin{matrix} R_3 \rightarrow R_3 - R_2 \end{matrix}$$

$$r(A) = 2$$

Consistency : A system of equations are said to be consistent if either they will have unique solution or many solutions and it to be inconsistent if they will have no solution.

$$2x + 3y = 8$$

$$x - 2y = 4$$

(unique solution)

$$x + 2y = 5$$

$$2x + 4y = 10$$

(many solutions)

$$x - y = 10$$

$$3x - 3y = 15$$

(No solution)

Consistency of a system of linear equations:-

Consider a system of m linear equations

$$\begin{aligned} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n &= b_1 \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n &= b_2 \\ \dots &\dots \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n &= b_m \end{aligned} \quad \dots (1)$$

Containing the unknowns $x_1, x_2, \dots, x_n$.

Writing the above equations in matrix form we get.

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}, B = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$$

$$C = \begin{bmatrix} A & B \end{bmatrix}$$

$$C = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} & b_1 \\ a_{21} & a_{22} & \dots & a_{2n} & b_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} & b_m \end{bmatrix}$$

A is the co-efficient matrix and

C is called augmented matrix

Rouche's Theorem: (Without proof)

The system of equations (1) is consistent if and only if the co-efficient matrix A and the augmented matrix C are of the same rank otherwise the system is inconsistent.

Procedure to test the consistency of a system of equations in n unknowns.

Find the rank of the co-efficient matrix A and the augmented matrix 'C' by reducing to the upper triangular form by elementary row operations.

(a) Consistent equations: If $\text{Rank } A = \text{Rank } C$

(i) Unique solution $\text{Rank } A = \text{Rank } C = n$

Where n = number of unknowns.

(ii) Infinite solution: $\text{Rank } A = \text{Rank } C = r, r < n$.

(b) Inconsistent equations if $\text{Rank } A \neq \text{Rank } C$

Example–2:

Show that the equations

$$2x + 6y = -11, 6x + 20y - 6z = -3, 6y - 18z = -1 \text{ are not consistent.}$$

Solution:

Writing the above equations in matrix form

$$\begin{bmatrix} 2 & 6 & 0 \\ 6 & 20 & -6 \\ 0 & 6 & -18 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -11 \\ -3 \\ -1 \end{bmatrix} \quad AX = B$$

$$\begin{aligned}
 & \begin{matrix} 2 & 6 & 0 \\ \hline 0 & 6 & -18 \\ 26 & 0 & -11 \end{matrix} \quad \begin{matrix} -11 \\ \hline -3 \end{matrix} \\
 & A = 620-6 \quad B = \begin{matrix} 0 & 6 & -18 \\ 26 & 0 & -11 \end{matrix} \quad C = [A:B] \\
 & \begin{matrix} 2 & 6 & 0 & -11 \\ \hline 0 & 6 & -18 & -1 \\ 26 & 0 & -11 & -3 \end{matrix} \\
 & C = \begin{matrix} 2 & 6 & 0 & -11 \\ \hline 0 & 6 & -18 & -1 \\ 26 & 0 & -11 & -3 \end{matrix} \quad \begin{matrix} 02 \\ \hline 6:30 \\ 0:11 \end{matrix} \quad \begin{matrix} R_2 - 3R_1 \\ R_3 - 2R_1 \end{matrix}
 \end{aligned}$$

The rank of C is 3

and rank of A is 2

Rank A $\neq$ Rank C.

\The system of equations are not consistent

Example–3:

Test consistency and solve:

$$5x + 3y + 7z = 4$$

$$3x + 2y + 2z = 9$$

$$7x + 2y + 10z = 5$$

Solution:

Writing the above equations in matrix form

$$\begin{aligned}
 & \begin{matrix} 5 & 3 & 7 \\ \hline 3 & 2 & 2 \\ 7 & 2 & 10 \end{matrix} \quad \begin{matrix} 4 \\ \hline 9 \\ 5 \end{matrix} \\
 & AX = B, C = [A:B] \\
 & \begin{matrix} 5 & 3 & 7 & 4 \\ \hline 3 & 2 & 2 & 9 \\ 7 & 2 & 10 & 5 \end{matrix} \\
 & C = \begin{matrix} 5 & 3 & 7 & 4 \\ \hline 3 & 2 & 2 & 9 \\ 7 & 2 & 10 & 5 \end{matrix} \quad \begin{matrix} R_2 - R_1 \\ R_3 - R_1 \end{matrix} \\
 & \begin{matrix} 5 & 3 & 7 & 4 \\ \hline 3 & 2 & 2 & 9 \\ 7 & 2 & 10 & 5 \end{matrix} \quad \begin{matrix} R_2 - R_1 \\ R_3 - R_1 \end{matrix} \\
 & \begin{matrix} 5 & 3 & 7 & 4 \\ \hline 3 & 2 & 2 & 9 \\ 7 & 2 & 10 & 5 \end{matrix} \quad \begin{matrix} R_2 - R_1 \\ R_3 - R_1 \end{matrix} \\
 & \begin{matrix} 5 & 3 & 7 & 4 \\ \hline 3 & 2 & 2 & 9 \\ 7 & 2 & 10 & 5 \end{matrix} \quad \begin{matrix} R_2 - R_1 \\ R_3 - R_1 \end{matrix}
 \end{aligned}$$

$$AX=B$$

$$C=[A:B]$$

$$C = \begin{bmatrix} 1 & 1 & 1 & 6 \\ 12 & 10 & 0 & 1 \\ 13 & m & 0 & 1 \\ 1 & 1 & 1 & 6 \end{bmatrix} \xrightarrow{R_2 - R_1} \begin{bmatrix} 1 & 1 & 1 & 6 \\ 0 & -1 & -1 & -5 \\ 0 & m-1 & -1 & -5 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & 1 & 1 & 6 \\ 0 & -1 & -1 & -5 \\ 0 & 0 & 0 & 0 \end{bmatrix} \xrightarrow{R_2 \times (-1)} \begin{bmatrix} 1 & 1 & 1 & 6 \\ 0 & 1 & 1 & 5 \\ 0 & 0 & 0 & 0 \end{bmatrix} \xrightarrow{R_1 - R_2} \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 5 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

- (i) There is no solution as $r(A) \neq r(C)$
i.e. $1-3=0$ or $1=3$ and $m-10=0$ or $m=10$
- (ii) There is a unique solution if $r(A) = r(C) = 3$
i.e., $1-3 \neq 0$ or $1 \neq 3$ and m has any value
- (iii) There are infinite solutions if $r(A) = r(C) = 2$ –
 $3 = 10$ or $1 = 3$ and $m-10 = 0$ or $m=10$

Assignments

1. Find the rank of the matrix $\begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 7 \\ 3 & 6 & 10 \end{bmatrix}$

2. Test the consistency & solve $4x$

$$-5y + z = 2$$

$$3x + y - 2z = 9$$

$$x + 4y + z = 5$$

3. Determine the values of a & b for which the system of equations $3x - 2y + z = b$

$$5x - 8y + 9z = 3$$

$$2x + y + az = -1$$

- (i) has a unique solution (ii) has no solution (iii) has infinite solutions.

COMPLEX NUMBERS

INTRODUCTION

We have the knowledge of integers, fractions and irrational number (all these constitute real numbers). But if we try to solve the equation $x^2 + 1 = 0$, we observe that these numbers are not adequate. Trying

to solve this equation, we arrive at $x^2 = -1$ i.e. $\sqrt{x} = -1$.

Square of a positive real number is positive and that of a negative real is also positive. So there is no real number whose square is negative. So we are to create a new kind of number. We define the square

root of a negative number as imaginary number' particularly $-1 = i$, the basic imaginary number.

Then $\sqrt{-4} = 2i$, $\sqrt{-2} = \sqrt{2}i$ and so on.

Imaginary numbers :

Taking $i = \sqrt{-1}$, we observe that

$$i^2 = -1$$

$$i^3 = -1.i = -i$$

$$i^4 = 1$$

Since $i^4 = 1$, $i = i^5 = i^9 = i^{13} = \dots = i^{4n+1}$, where n is an integer.

$$i^6 = i^{10} = i^{14} = \dots = i^{4n+2}$$

$$i^3 = i^7 = i^{11} = i^{15} = \dots = i^{4n+3}$$

$$i^4 = i^8 = i^{12} = i^{16} = \dots = i^{4n}$$

COMPLEX NUMBERS

The numbers of the form $a + ib$ where a and b are real numbers and $i = \sqrt{-1}$ are known as complex numbers.

In complex number $z = a + ib$, the real numbers a and b are respectively known as real and imaginary parts of z and we write :

$\text{Re}(z) = a$ and $\text{Im}(z) = b$

Thus the set C of all complex numbers is given by $C = \{z : z = a + ib, \text{ where } a, b \in \mathbb{R}\}$

Purely real and purely imaginary numbers :

A complex number z is said to be

(i) Purely real, if $\text{Im}(z) = 0$

(ii) Purely imaginary, if $\text{Re}(z) = 0$

Thus, $2, -7, \sqrt{3}$ etc are all purely real numbers.

While $2i, i\sqrt{3}, \frac{-1}{2}i$ etc are purely imaginary.

Conjugate of a complex number :

The conjugate of a complex number ' z ', denoted by $\bar{z}$ is the complex number obtained by changing the sign of imaginary part of z .

$$\text{e.g. } \overline{2 + 3i} = (2 - 3i); \quad \overline{3 + 5i} = (3 - 5i),$$

$$\overline{6i} = -6i; \quad \overline{-2i} = 2i$$

Modulus of a complex number : If $z = x + iy$ be a complex number, the modulus of z , written as $|z|$ is a real number $\sqrt{x^2 + y^2}$.

For $z = 3 + 4i$, $|z| = \sqrt{3^2 + 4^2} = 5$.

Also $|\bar{z}| = |z|$.

If $z = x + iy$, $\bar{z} = x - iy$.

$$|z| = \sqrt{x^2 + y^2}, |\bar{z}| = \sqrt{x^2 + (-y)^2} = \sqrt{x^2 + y^2}$$

SUM DIFFERENCE AND PRODUCT OF COMPLEX NUMBERS

For any complex number

$$z_1 = (a + ib) \text{ and } z_2 = (c + id)$$

we define

$$(i) \ z_1 + z_2 = (a + ib) + (c + id) = [(a + c) + i(b + d)]$$

$$(ii) \ z_1 - z_2 = (a + ib) - (c + id) = [(a - c) + i(b - d)]$$

$$(iii) \ z_1 z_2 = (a + ib)(c + id) = [(ac - bd) + i(ad + bc)]$$

CUBE ROOTS OF UNITY

Let $\sqrt[3]{1} = x$, then

$$x^3 = 1 \text{ [on cubing both sides]}$$

$$\Rightarrow x^3 - 1 = 0 \quad \Rightarrow (x - 1)(x^2 + x + 1) = 0$$

$$\Rightarrow x - 1 = 0 \quad \text{or} \quad x^2 + x + 1 = 0$$

$$\Rightarrow x = 1 \quad \text{or} \quad x = \frac{-1 \pm \sqrt{1-4}}{2}$$

$$\Rightarrow x = 1 \quad \text{or} \quad x = \frac{-1 \pm i\sqrt{3}}{2}$$

$$\therefore \text{The cube roots of unity are } 1, \frac{-1 + i\sqrt{3}}{2} \text{ and } \frac{-1 - i\sqrt{3}}{2}$$

Clearly one of the cube roots of unity is real and the other two are complex.

Example - 1 : Express in the form $a + ib$

$$\begin{aligned} (i) \quad & \frac{3+5i}{2-3i} \quad (ii) \quad \frac{(1+i)^2}{3-i} \\ \text{Sol}^n : (i) \quad & \frac{3+5i}{2-3i} = \frac{(3+5i)(2+3i)}{(2-3i)(2+3i)} = \frac{6+10i+9i+15i^2}{4-9i^2} = \frac{-9+19i}{13} = \frac{-9}{13} + \frac{19}{13}i \\ (ii) \quad & \frac{(1+i)^2}{3-i} = \frac{(1+i^2+2i)(3+i)}{(3-i)(3+i)} = \frac{6i-2i^2}{9-i^2} = \frac{6i+2}{10} = \frac{13}{5} + \frac{3}{5}i \end{aligned}$$

Example - 2 : Find the value of $i^{17} + i^{20} - i^{13}$

$$\begin{aligned} \text{Sol}^n : i^{17} + i^{20} - i^{13} &= i^{16} \cdot i + i^{20} - i^{12} \cdot i = (i^2)^8 \cdot i + (i^2)^{10} - (i^2)^6 \cdot i \\ &= (-1)^8 i + (-1)^{10} - (-1)^6 i = i + 1 - i = 1 \end{aligned}$$

Example - 3 : If $1, w, w^2$ are the cube roots of unity prove that

$$(a) \quad (1-w)(1-w^2)(1-w^4)(1-w^5) = 9$$

$$\begin{aligned} \text{Sol}^n : \text{L.H.S. } (1-w)(1-w^2)(1-w^4)(1-w^5) \\ &= (1-w)(1-w^2)(1-w^3 \cdot w)(1-w^3w^2) \\ &= (1-w)(1-w^2)(1-w)(1-w^2) \\ &= (1-w)^2(1-w^2)^2 = [(1-w)(1-w^2)]^2 \end{aligned}$$

$$= [(1 - w - w^2 + w^3)]^2 = (2 - w - w^2)^2$$

$$= (2+1)^2 = 3^2 = 9$$

Example-4: Find square roots of

(a) $3 + 4i$

Solⁿ : (a) Let $x, y \in \mathbb{R}$, $x + iy = \sqrt{3 + 4i}$

$$x^2 - y^2 + i 2xy = 3 + 4i$$

Equating real and imaginary parts

$$x^2 - y^2 = 3 \text{ and } 2xy = 4$$

$$(x^2 + y^2)^2 = (x^2 - y^2)^2 + 4x^2y^2 = 25$$

Hence $x^2 + y^2 = \pm 5$, But since $x^2 + y^2$ is non-negative, we have

$$x^2 + y^2 = 5$$

$$x^2 - y^2 = 3$$

$$2x^2 = 8$$

$$\text{i.e., } x^2 = 4, \text{ i.e., } x = \pm 2, y^2 = 1 \text{ i.e., } y = \pm 1$$

Hence square roots of $3 + 4i = \pm (2 + i)$

Assignment

1. If w be the cube roots of unity, then prove that $(1 - w + w^2)^7 + (1 + w + w^2)^7 = 128$
2. Find square roots of $-5 + 12\sqrt{-1}$

LINEAR DIFFERENTIAL EQUATIONS

Introduction:

The Mathematical formulation of many problems in science, Engineering and Economics gives rise to differential Equations.

For example: The problem of motion of a satellite

- 1 The flow of fluids.
- 1 The flow of current in an electric circuit
- 1 The growth of population
- 1 The Conduction of heat in rod etc lead to differential equations

Definition of Differential Equation:

A differential equation is an equation involving derivatives of one or more dependent variables with respect to one or more independent variables.

There are two types of Differential Equation

1. Ordinary differential Equation
2. Partial differential Equation

Example:

- (a) $\frac{dy}{dx} + y = x^2$
- (b) $\frac{d^3y}{dx^3} + 3\frac{d^2y}{dx^2} + 2\frac{dy}{dx} + y = 0$
- (c) $\frac{\partial u}{\partial t} + \frac{\partial^2 u}{\partial x^2} = 4$

Linear differential Equation:

Linear differential Equations are those in which the dependent variable and its derivatives occur only in the first degree and are not multiplied together.

The differential Equation of the form

$$\frac{d^n y}{dx^n} + k_1 \frac{d^{n-1} y}{dx^{n-1}} + k_2 \frac{d^{n-2} y}{dx^{n-2}} + \dots + k_n y = X \dots \dots \dots (1)$$

Is known as linear differential equation with constant coefficients. Where $k_1, k_2, \dots, k_n$ are constant, X is the function of x .

There are two types of linear differential equation

1. Homogeneous LDE
2. Non Homogeneous LDE

Homogeneous Linear Differential Equation:

If R.H.S. of Equation (1) is equal to zero then we get homogeneous LDE.

$$\text{ie } \frac{d^n y}{dx^n} + k_1 \frac{d^{n-1} y}{dx^{n-1}} + k_2 \frac{d^{n-2} y}{dx^{n-2}} + \dots + k_n y = 0$$

Where $f(x)$ is the function of ' x '

The general solution format of Equation (1) of the form $(C.S = C.F + P.I)$

Where C.S. – Complete Solution

C.F – Complementary function

P.I – Particular integral

So complete solution of Equation becomes $(y = C.F + P.I)$

Note -1: In case of Homogeneous LDE

$$C.S = C.F \text{ [where } P.I = 0]$$

Note -2: In case of Non-Homogeneous LDE

$$C.S = C.F + P.I$$

Operator:

Denoting $\frac{d}{dx}, \frac{d^2}{dx^2}, \frac{d^3}{dx^3}, \dots$ by $D, D^2, D^3, \dots$

So that $\frac{dy}{dx} = Dy$

$$\frac{d^2 y}{dx^2} = D^2 y$$

.....

$$\frac{d^n y}{dx^n} = D^n y$$

Where D – Derivative

Then $\frac{1}{D}$ – Integration

Then operator form of equation (1)

$$\text{becomes } D^n y + k_1 D^{n-1} y + k_2 D^{n-2} y + \dots + k_n y = X$$

$$(D^n + k_1 D^{n-1} + k_2 D^{n-2} + \dots + k_n) y = X$$

$$F(D)y = X \dots \dots$$

(2)

$$\text{Where } F(D) = D^n + k_1 D^{n-1} + k_2 D^{n-2} + \dots + k_n \text{ of function } D$$

Auxiliary Equation(AE)

Putting the coefficient of y equal to Zero in Equation (2) we get an Auxiliary Equation.

i.e. $F(D)=0$

i.e. $D^n + k_1 D^{n-1} + k_2 D^{n-2} + \dots + k_n = 0$

Depending on the value of 'D' in Auxiliary Equation, complementary functions are of different types.

Case-I: If roots are real & Different

Let m_1 & m_2 are two real roots and different

i.e. $m_1 \neq m_2$

Then C.F = $C_1 e^{m_1 x} + C_2 e^{m_2 x}$

Where C_1, C_2 , are arbitrary constants

Case-II: If roots are real & Equal

Let m_1 & m_2 are two real roots & Equal

i.e. $m_1 = m_2$

The C.F = $(C_1 + C_2 x) e^{m_1 x}$

Similarly if $m_1 = m_2 = m_3$ (Three roots are Equal)

Then C.F = $(C_1 + C_2 x + C_3 x^2) e^{m_1 x}$

Case-III: If roots are Complex conjugate

Let $m_1 = a \pm ib$ are conjugate complex roots

Then C.F = $e^{ax} \{C_1 \cos bx + C_2 \sin bx\}$

Case-IV: If two conjugate complex roots are equal

Let $m_1 = m_2 = a \pm ib$ are equal

Then C.F = $e^{ax} \{C_1 + C_2 x\} \cos bx + (C_3 + C_4 x) \sin bx$

Example-1:

Solve $\frac{d^2 y}{dx^2} - 8 \frac{dy}{dx} + 15y = 0 \dots\dots\dots (1)$

Solution:

The operator from equation (1) becomes

$(D^2 - 8D + 15)y = 0$

So Auxiliary Equation $D^2 -$

$8D + 15 = 0$

$\therefore (D-3)(D-5) = 0$

$\therefore D = 3, 5$

Then C.F = $C_1 e^{3x} + C_2 e^{5x}$

So complete Solution

$y = C_1 e^{3x} + C_2 e^{5x} \quad (\text{Ans})$

Example–2:

$$\text{Solve } \frac{d^2 y}{dx^2} - 6 \frac{dy}{dx} + 9y = 0$$

Solution:

The operator from of given equation is

$$(D^2 - 6D + 9)y = 0$$

$$\text{Then A.E } D^2 - 6D + 9 = 0$$

$$D(D-3)^2 = 0$$

$$D = 3, 3$$

$$C.F = (C_1 + C_2 x)e^{3x}$$

$$\text{Then C.Sy} = (C_1 + C_2 x)e^{3x} \quad (\text{Ans})$$

Example–3:

$$\text{Solve } (D^2 + 4D + 5)y = 0$$

Solution:

$$\text{So A.E } D^2 + 4D + 5 = 0$$

$$D = \frac{-4 \pm \sqrt{16 - 4 \cdot 1 \cdot 5}}{2 \cdot 1}$$

$$= \frac{-4 \pm \sqrt{-4}}{2}$$

$$= \frac{-4 \pm 2i}{2} = -2 \pm i$$

$$\text{Then C.F} = e^{-2x} \{ C_1 \cos x + C_2 \sin x \}$$

$$\text{So C.Sy} = e^{-2x} \{ C_1 \cos x + C_2 \sin x \} \quad (\text{Ans})$$

Procedure to finding particular Integral.

We know that $F(D)y = X$

$$y = \frac{X}{F(D)}$$

Depending upon nature of 'X', Particular integral are different types

Case –1 : When $X = e^{ax}$

$$\text{Then P.I} = \frac{e^{ax}}{F(a)} \text{ where } D = a$$

$$\text{If } F(a) \neq 0, \text{ Then PI} = \frac{X e^{ax}}{F'(a)} \text{ provided } F(a) \neq 0$$

$$\text{If } F(a) = 0, \text{ Then PI} = \frac{X^2 e^{ax}}{F''(a)} \text{ provided } F''(a) \neq 0$$

And soon.

Case–2: When $X = \sin(ax+b)$ or $\cos(ax+b)$

$$\text{Then PI} = \frac{\sin(ax+b)}{FD^2} \text{ Put } D^2 = -a^2$$

But not $D = -a$

$$\frac{\sin(ax+b)}{FD^2} = \frac{\sin(ax+b)}{F(-a^2)} \text{ provided } F(-a^2) \neq 0$$

If $F(-a^2) = 0$, The above rule fails & we proceed further

$$\text{ie P.I} = \frac{x \sin(ax+b)}{F(-a^2)}, \text{ Provided } F(-a^2) \neq 0$$

$$\text{If } F(-a^2) = 0, \text{ Then P.I} = \frac{\sin(ax+b)}{F(-a^2)}, \text{ Provided } F(-a^2) \neq 0$$

And soon

Case–3: When $X = e^{ax}v$, Where $v = \text{function of } x$

$$\text{Then PI} = \frac{e^{ax}v}{FD}$$

$$= \frac{1}{F(D+a)} e^{ax}v$$

Similarly when $X = e^{-ax}v$

$$\text{Then PI} = e^{-ax} \left(\frac{1}{FD+a} \right) v$$

Case–4: When $X = x^m$ (ie, $x, x^2, x^3, \dots$)

$$\text{Then PI} = \frac{x^m}{FD(D+1)(D+2)\dots(D+m-1)}$$

Convert $F(D)$ into $\{1+F(D)\}$ or $\{1-F(D)\}$ by taking D^m (if possible). Then by using Binomial Theorem we find solution.

Case–5: When $X = xv$

$$\text{Then P.I} = \frac{xv}{FD} = \frac{x}{F(D)} \frac{v}{D}$$

Where $F(D)$ is the Derivative of $F(D)$

Case–6: When x is any other function

$$\text{Then P.I.} = \frac{x}{F(D)}$$

Convert $F(D)$ into $(D - a)$ or $(D + a)$ factor form

$$\text{Then if } \frac{x}{D - \alpha} = e^{\alpha x} \int x e^{-\alpha x} dx \quad \text{if } \frac{x}{D - \alpha} = e^{-\alpha x} \int x e^{\alpha x} dx$$

Example–4:

$$\text{Find P.I. of } (D^2 + 6D + 3)y = e^{2x}$$

Solution:

$$\text{P.I.} = \frac{e^{2x}}{D^2 + 6D + 3} \quad \text{put } D = a$$

$$\text{i.e. } D = 2$$

$$\text{Then P.I.} = \frac{e^{2x}}{(2)^2 + 6(2) + 3}$$

$$= \frac{e^{2x}}{4 + 12 + 3} = \frac{e^{2x}}{19} \quad (\text{Ans})$$

Example–5:

$$\text{Solve } \frac{d^3 y}{dx^3} - 3 \frac{d^2 y}{dx^2} + 4 \frac{dy}{dx} - 2y = e^x + \cos x$$

Solution:

The operator form of given equation becomes $(D^3 - 3D^2 + 4D - 2)y = e^x + \cos x$

$$- 3D^2 + 4D - 2) y = e^x + \cos x$$

$$\text{So A.E. } D^3 - 3D^2 + 4D - 2 = 0$$

$$D - 1, 1 \pm i$$

$$D = 1, 1 \pm i$$

$$\text{C.F.} = C_1 e^x + e^x \{ C_2 \cos x + C_3 \sin x \}$$

$$\text{Then P.I.} = \frac{e^x + \cos x}{D^3 - 3D^2 + 4D - 2}$$

$$\frac{e^x}{D^3 - 3D^2 + 4D - 2} + \frac{\cos x}{D^3 - 3D^2 + 4D - 2}$$

$$= (D - 1)(D^2 - 2D + 2) + D^3 - 3D^2 + 4D - 2$$

$$\frac{e^x}{D - 1 - 2 + 2} + \frac{\cos x}{+1D - 3 - 1 + 4D - 2}$$

Example–7:

Solve $\frac{d^2 y}{dx^2} + 9y = x \cos x$

Solution:

The operator form is $(D^2 + 9) y = x \cos x$

So A.E. $D^2 + 9 = 0$

$D^2 = -9$

$D = \pm 3i$

$D = \pm 3i$

C.F. $= C_1 \cos 3x + C_2 \sin 3x$

Now P.I. $= \frac{x \cos x}{D^2 + 9}$

Here $F(D) = D^2 + 9$

$F'(D) = 2D$

Then P.I. $= \frac{F(D)}{F'(D)} \cdot \frac{1}{D} \cdot x \cos x$

$= \frac{1}{D} \cdot \frac{2D \cos x}{D^2 + 9} \cdot x \cos x$ put $D^2 = -1$

$= \frac{1}{D} \cdot \frac{2D \cos x}{D^2 + 9} \cdot x \cos x$

$= \frac{1}{D} \cdot \frac{2D \cos x}{D^2 + 9} \cdot x \cos x$

$= \frac{1}{D} \cdot \frac{2D \cos x}{D^2 + 9} \cdot x \cos x$

$= \frac{1}{D} \cdot \frac{2D \cos x}{D^2 + 9} \cdot x \cos x$

$= \frac{1}{D} \cdot \frac{2D \cos x}{D^2 + 9} \cdot x \cos x$

So C.Sy. $= C_1 \cos x + C_2 \sin x + \frac{4x \cos x + \sin x}{32}$

(Ans)

32

Example–8:

Solve $\frac{d^2 y}{dx^2} + 4y = x^2$

Solution:

The operation form given equation becomes

$(D^2 + 4) y = x^2$

So A.E. $D^2 + 4 = 0$

$$D^2 = -4$$

$D = \sqrt{-4}$

$D = \pm 2i$

$$C.F. = C_1 \cos 2x + C_2 \sin 2x$$

Then P.I. = $\frac{x^2}{D^2 + 4}$

$$= \frac{1}{D^2 + 4} x^2$$

$$= \frac{1}{D^2 + 4} x^2 = \frac{1}{D^2 + 4} x^2$$

by using Binomial theorem

$$= \frac{1}{D^2 + 4} x^2 = \frac{1}{D^2 + 4} x^2$$

$$= \frac{1}{D^2 + 4} x^2 = \frac{1}{D^2 + 4} x^2$$

So C.F. = $C_1 \cos 2x + C_2 \sin 2x + \frac{x^2}{8} - 1$ (Ans)

Other Method for finding P.I.:

Method of variation of Parameters:

This method is applied to equations of the form

$$y'' + py' + qy = x$$

Where p, q & x are functions of x .

Then P.I. = $\int y_1 \frac{y_2 x dx}{W} + \int y_2 \frac{y_1 x dx}{W}$

Where y_1 & y_2 are the solutions of $y'' + py' + qy = 0$ of the form $y = c_1 y_1 + c_2 y_2$ & W is called the wronskian of y_1 & y_2

Calculate by formula $W(y_1, y_2) = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$

Example–9:

$$\text{Solve } \frac{d^2 y}{dx^2} + y = \operatorname{cosec} x$$

Solution:

The operator form of given equation is

$$(D^2 + 1)y = \operatorname{cosec} x$$

$$\text{So A.E } D^2 + 1 = 0$$

$$\text{P } D^2 = -1$$

$$\text{P.D. } = \sqrt{-1} = 0 \pm i$$

$$\text{C.F. } = C_1 \cos x + C_2 \sin x$$

$$\text{Here } y_1 = \cos x, y_2 = \sin x$$

$$W(y_1, y_2) = \begin{vmatrix} \cos x & \sin x \\ -\sin x & \cos x \end{vmatrix}$$

$$= \cos^2 x + \sin^2 x = 1$$

$$\begin{aligned} \text{Then P.I.} &= -\cos x \int \frac{\sin x \cdot \operatorname{cosec} x}{1} dx + \sin x \int \frac{\sin x \cdot \operatorname{cosec} x}{1} dx \\ &= -\cos x \int \sin x \cdot \frac{1}{\sin x} dx + \sin x \int \cos x \cdot \frac{1}{\sin x} dx \\ &= -\cos x \int dx + \sin x \int \cot x dx \\ &= -\cos x(x) + \sin x \ln \sin x \end{aligned}$$

$$\text{So C.Sy} = C_1 \cos x + C_2 \sin x + \sin x \ln \sin x - x \cos x \text{ (Ans.)}$$

Partial Differential Equation

Let $z = f(x, y)$ be a function containing two independent variables x & y and z is the Dependent variable.

Notation: Let $z = f(x, y)$ be a function of x & y

$$\begin{aligned} \text{Then } \frac{\partial z}{\partial x} &= p & \frac{\partial z}{\partial y} &= q \\ \frac{\partial^2 z}{\partial x^2} &= r & \frac{\partial^2 z}{\partial y^2} &= t \\ \frac{\partial^2 z}{\partial x \partial y} &= s \end{aligned}$$

Formation of Partial Differential Equation

A partial differential equation can be formed by

- (i) Eliminating arbitrary constant.
- (ii) Eliminating arbitrary function.

Example–10:

Form a partial differential equation by eliminating function

$$Z = f(x^2 + y^2) \quad \dots(1)$$

Solution:

Differentiating partially w.r.t. x & y in equation (1) we get

$$\frac{\partial Z}{\partial x} = f'(x^2 + y^2) \cdot 2x \quad (\text{taking } y \text{ as a constant})$$

$$p = f'(x^2 + y^2) \cdot 2x \quad \dots(2)$$

$$\text{Similarly } q = f'(x^2 + y^2) \cdot 2y \quad \dots(3)$$

$$\text{Dividing (2) \& (3) we get} \quad \frac{p}{q} = \frac{f'(x^2 + y^2) \cdot 2x}{f'(x^2 + y^2) \cdot 2y}$$

$$\frac{p}{q} = \frac{x}{y} \quad \dots(4)$$

$$py - qx = 0 \quad (\text{Ans.})$$

Linear Equation of the First order:

A Linear partial differential equation of the 1st order is of the form

$$Pp + Qq = R$$

Where P, Q & R are functions of x, y, z.

This equation is also known as Lagrange's Linear equation

NOTE:

The general solution of the linear partial differential equation $Pp + Qq = R$ is $f(a, b) = 0$

$$\text{Or } a = f(b)$$

$$\text{Or } b = f(a)$$

Where f is an arbitrary function & $u(x, y, z) = a$ & $v(x, y, z) = b$ form the solution of the equation

$$\frac{dx}{P} = \frac{dy}{Q} = \frac{dz}{R}$$

Then that can be solved by two methods

- (1) Method for Grouping
- (2) Method for Multipliers

Method of grouping:

Take any two fractions from Subsidiary Equation such that the 3rd variable is absent or it may be cancelled.

$$\text{For example take } \frac{dx}{P} = \frac{dy}{Q} \quad (\text{such that } z \text{ may be absent})$$

After Integration we get $f(x, y) = a$

Similarly we take $\frac{dy}{dz} = \frac{dz}{dz}$

R

After Integration $f(y, z) = b$

So general solution is $a = f(b)$

or $b = f(a)$

or $f(a, b) = 0$

Method for Multipliers

Let us choose the multiplier's (P , Q , R) such

That $PP' + QQ' + RR' = 0$

Then we write $P'dx + Q'dy + R'dz = 0$

On Integration we get $f(x, y, z) = a$

Similarly choosing the multipliers (P^2, Q^2, R^2) such that

$PP^2 + QQ^2 + RR^2 = 0$

On Integration we get $g(x, y, z) = b$

So general solution is $a = f(b)$ or $f(a, b) = 0$

Example–11:

Solve $y^2z^p + z^2x^q = y^2x$

Solution:

It is of the form $Pp + Qq = R$

Where $P = y^2z$, $Q = z^2x$, $R = y^2x$

$$\text{So it is } P \frac{dx}{yz} = \frac{dy}{zx} = \frac{dz}{yx}$$

Taking 1st and 3rd fraction, we get

$$\frac{dx}{yz} = \frac{dz}{yx} \quad (\text{Here } 3^{\text{rd}} \text{ variable } y^2 \text{ is cancelled})$$

$$\Rightarrow xdx = zdz$$

Integrating both sides we get $\int xdx = \int zdz$

$$\Rightarrow \frac{x^2}{2} = \frac{z^2}{2} + C$$

$$\Rightarrow x^2 - z^2 = 2C = a$$

Similarly taking 2nd and 3rd

$$\text{i.e. } \frac{dy}{z^2x} = \frac{dz}{y^2x}$$

$$\Rightarrow y^2dy = z^2dz$$

Integrating both sides we get

$$\int \frac{y^3 - z^3}{33} = \int \frac{1}{33} + C^1$$

$$\int y^3 - z^3 = 3C^1 = b$$

$$\text{So general solution is } x^2 - z^2 = f(y^3 - z^3) \quad (\text{Ans.})$$

Example-12:

$$\text{Solve } x(z^2 - y^2)p + y(x^2 + y^2)q = z(y^2 - x^2)$$

Solution:

It is the equation of the form

$$Pp + Qq = R$$

$$\text{Where } P = x(z^2 - y^2), Q = y(x^2 - z^2), R = z(y^2 - x^2)$$

$$\text{So it is } \frac{dx}{x(z^2 - y^2)} = \frac{dy}{y(x^2 - z^2)} = \frac{dz}{z(y^2 - x^2)}$$

Let us choose multipliers (x, y, z) . i.e. $P' = x, Q' = y, R' = z$

$$= z \text{ Such that } x \cdot x(z^2 - y^2) + y \cdot y(x^2 - z^2) + z \cdot z(y^2 - x^2)$$

$$= x^2 z^2 - x^2 y^2 + y^2 x^2 - y^2 z^2 + z^2 y^2 - z^2 x^2$$

$$= 0$$

Then we write $x dx + y dy + z dz = 0$

On integration we set

$$\frac{x^2}{2} + \frac{y^2}{2} + \frac{z^2}{2} = c$$

$$\int x^2 + y^2 + z^2 = 2C = a$$

$$\text{Again choose the multipliers } \frac{1}{x}, -\frac{1}{y}, \frac{1}{z} \text{ i.e. } P^2 = \frac{1}{x}, Q^2 = -\frac{1}{y}, R^2 = \frac{1}{z}$$

$$\text{Such that } \frac{1}{x}(z^2 - y^2) - \frac{1}{y}(x^2 - z^2) + \frac{1}{z}(y^2 - x^2)$$

$$= z^2 - y^2 + x^2 - z^2 + y^2 - x^2 = 0$$

$$\text{Then } \frac{1}{x} dx - \frac{1}{y} dy + \frac{1}{z} dz = 0$$

On integration we get

$$\log x + \log y + \log z = \log b$$

$$\int \log(xyz) = \log b$$

$$\int xyz = b$$

$$\text{So general solution is } x^2 + y^2 + z^2 = f(xyz) \quad (\text{Ans})$$

Assignment

Solve the followings:

1. $\frac{d^2y}{dx^2} + 3 \frac{dy}{dx} - 10y = 6e^{4x}$

2. $y'' + 3y' + 2y = 4\cos^2 x$

3. $\frac{d^2y}{dx^2} - 2 \frac{dy}{dx} + y = x^2 e^x$

4. $(D^2 + a^2)y = k \cos(ax + b)$

5. $(D - 2)^2 y = 8(e^{2x} + \sin 2x)$

6. $\frac{d^3y}{dx^3} - \frac{d^2y}{dx^2} - 6 \frac{dy}{dx} = 1 + x^2$

CHAPTER–3**LAPLACE TRANSFORMS****GAMMA FUNCTION:**

The gamma function is defined as

$$\Gamma(n) = \int_0^{\infty} e^{-x} x^{n-1} dx, n > 0 \quad \dots(1)$$

It defines a function of n for positive values of n .

Value of $\Gamma(1)$:

We have,

$$\Gamma(1) = \int_0^{\infty} e^{-x} x^0 dx = \int_0^{\infty} e^{-x} dx = \left[-e^{-x} \right]_0^{\infty} = 1$$

$$\text{Hence, } \Gamma(1) = 1 \quad \dots(2)$$

Reduction formula for $\Gamma(n)$: We

have,

$$\begin{aligned} \Gamma(n+1) &= \int_0^{\infty} e^{-x} x^n dx \quad [\text{Integrating by parts}] \\ &= \left[-x^n e^{-x} \right]_0^{\infty} + n \int_0^{\infty} e^{-x} x^{n-1} dx = 0 + n\Gamma(n) \end{aligned}$$


$$\Gamma(n+1) = n\Gamma(n), \quad \dots(3)$$

which is the reduction formula for $\Gamma(n)$.

Using the reduction formula for $\Gamma(n)$, we can write the value of $\Gamma(n)$ in the form,

$$\Gamma(n) = \frac{\Gamma(n+1)}{n} \dots \dots \dots (4)$$

Thus (1) and (4) together give a complete definition of $\Gamma(n)$ defined for all values of n except when n is zero or a negative integer and its graph is as shown in the following figure.



VALUE OF $G(n)$ IN TERMS OF FACTORIAL

Using $G(n+1) = nG(n)$ successively, we get

$$G(2) = G(1+1) = 1 \times G(1) = 1!$$

$$G(3) = G(2+1) = 2 \times G(2) = 2 \times 1 = 2!$$

$$G(4) = G(3+1) = 3 \times G(3) = 3 \times 2! = 3!$$

.....

In general $G(n+1) = n!$, provided n is a positive integer.

Taking $n = 0$, it defines $0! = G(1) = 1$

Thus, $G(n+1) = n!$ (for $n = 0, 1, 2, 3, \dots$).....(5)

Value of G^{-1}

2:

We have,

$$G^{-1} = \frac{1}{G} = \frac{1}{n!} = \frac{1}{n(n-1)(n-2)\dots 1} = \frac{1}{n!}$$

$$= 2 \int_0^{\infty} e^{-y^2} dy, \text{ Which is also } = 2 \int_0^{\infty} e^{-x^2} dx$$

$$\int_0^{\infty} \int_0^{\infty} e^{-(x^2+y^2)} dx dy = \int_0^{\infty} \int_0^{\infty} e^{-r^2} r dr d\theta = \int_0^{\infty} \int_0^{2\pi} e^{-r^2} r dr d\theta$$

$$= \int_0^{\infty} e^{-r^2} r dr \int_0^{2\pi} d\theta = \int_0^{\infty} e^{-r^2} r dr \cdot 2\pi$$

$$= 2\pi \int_0^{\infty} e^{-r^2} r dr = 2\pi \left[-\frac{1}{2} e^{-r^2} \right]_0^{\infty} = 2\pi \left[-\frac{1}{2} e^{-\infty} + \frac{1}{2} e^{-0} \right] = 2\pi \left[0 + \frac{1}{2} \right] = \pi$$

Hence $G_{\zeta} = \sqrt{p} = 1.772 \dots \dots \dots (6)$

Example-1:

$\mathfrak{a}3\mathfrak{o} \quad \mathfrak{a}5\mathfrak{o} \quad \mathfrak{a}1\mathfrak{o}$
 $G_5 \div \times G_5 \div \times G_5 \div$
 Evaluate, $\mathfrak{e}2\mathfrak{o} \quad \mathfrak{e}2 \mathfrak{o} \quad \mathfrak{e}2\mathfrak{o}$
 $G(4).G(6).G(8)$

Solution:

$$\frac{\begin{array}{ccc} \mathfrak{a}3\mathfrak{o} & \mathfrak{a}5\mathfrak{o} & \mathfrak{a}1\mathfrak{o} \\ \text{G}\mathfrak{c}\text{---}\div\times\text{G}\mathfrak{c}\div & \times\text{G} & \text{---}\div \\ \text{We have,}\mathfrak{e}2\mathfrak{o} & \mathfrak{e}2\mathfrak{o} & \mathfrak{e}2\mathfrak{o} \end{array}}{\text{G}(4).\text{G}(6).\text{G}(8)}$$

[illegible]

Example-2:

EvaluateG(-3.5)

Solution:

$$G(n+1)$$

We know that, $G(n) = \frac{n(n+1)}{2}$

For all n expect n is zero or a negative integer.

Now, we have,

$$\begin{aligned} G(-3.5) &= \frac{G(-3.5+1)}{-3.5} = \frac{G(-2.5)}{-3.5} = \frac{G(-2.5+1)}{(-3.5)(-2.5)} = \frac{G(-1.5)}{(3.5)(2.5)} \\ &= \frac{G(-1.5+1)}{(3.5)(2.5)(-1.5)} = \frac{G(-0.5)}{(3.5)(2.5)(-1.5)} = \frac{G(-0.5+1)}{(3.5)(2.5)(-1.5)(-0.5)} \\ &= \frac{G(0.5)}{(3.5)(2.5)(1.5)(0.5)} = \frac{\sqrt{p}}{(3.5)(2.5)(1.5)(0.5)} = 0.27 \end{aligned}$$

$$\backslash G(-3.5)=0.27$$

Laplace transforms :

Definition:

Let $f(t)$ be a function of t defined for all positive values of t . Then the Laplace transform of $f(t)$, denoted by $L\{f(t)\}$ is defined by

$$L\{f(t)\} = \int_0^{\infty} e^{-st} f(t) dt$$

Provided that the integral exists. s is a parameter which may be a real or complex number.

$L\{f(t)\}$ being clearly a function of s is briefly written as $f(s)$.

i.e., $L\{f(t)\} = f(s)$.

This implies that, $f(t) = L^{-1}\{f(s)\}$

Then $f(t)$ is called the inverse Laplace transform of $f(s)$.

The symbol L , which transforms $f(t)$ into $f(s)$, is called the Laplace transformation operator.

CONDITIONS FOR THE EXISTENCE:

The Laplace transform of $f(t)$ i.e., $\int_0^{\infty} e^{-st} f(t) dt$ exists for $s > a$, if

- (i) $f(t)$ is continuous
- and (ii) $\lim_{t \rightarrow \infty} \{e^{-at} f(t)\}$ is finite.

TRANSFORMS OF ELEMENTARY FUNCTIONS:

The direct application of the definition gives the following formulae: 1

$$(1) L\{1\} = \frac{1}{s} \quad (s > 0)$$

$$(2) L\{t^n\} = \frac{n!}{s^{n+1}}, \text{ when } n=0, 1, 2, 3, \dots$$

$$= \frac{1}{s^{n+1}}, \text{ otherwise } (s > 0)$$

$$(3) L\{e^{at}\} = \frac{1}{s-a} \quad (s > a)$$

$$(4) L\{\sin at\} = \frac{a}{s^2 + a^2} \quad (s > 0)$$

$$(5) L\{\cos at\} = \frac{s}{s^2 - a^2} \quad (s > 0)$$

$$(6) L\{\sinh at\} = \frac{a}{s^2 - a^2} \quad (s > |a|)$$

$$(7) \quad L\{\cosh at\} = \frac{s}{s^2 - a^2} \quad (s > |a|)$$

PROOFS:

$$(1) \quad L\{1\} = \int_0^{\infty} e^{-st} \cdot 1 \cdot dt = \left[-\frac{e^{-st}}{s} \right]_0^{\infty} = \frac{1}{s}, \text{ if } s > 0$$

$$(2) \quad L\{t^n\} = \int_0^{\infty} e^{-st} \cdot t^n \cdot dt = \int_0^{\infty} e^{-pt} \cdot t^n \cdot dt \quad \text{on putting } st = p$$

$$= \frac{n!}{s^{n+1}}, \text{ if } n > -1 \text{ and } s > 0$$

s^{n+1}

If n is a positive integer, $G(n+1) = n!$. Therefore, $L\{t^n\} = \frac{n!}{s^{n+1}}, \text{ if } s > 0$

$$(3) \quad L\{e^{at}\} = \int_0^{\infty} e^{-st} \cdot e^{at} dt = \int_0^{\infty} e^{-(s-a)t} dt = \left[\frac{e^{-(s-a)t}}{-(s-a)} \right]_0^{\infty} = \frac{1}{s-a}, \text{ if } s > a$$

$$(4) \quad L\{\sin at\} = \int_0^{\infty} e^{-st} \sin at dt = \left[\frac{e^{-st}}{s^2 + a^2} (-s \sin at - a \cos at) \right]_0^{\infty} = \frac{a}{s^2 + a^2}, \text{ if } s > 0$$

$$(5) \quad L\{\cos at\} = \int_0^{\infty} e^{-st} \cos at dt = \left[\frac{e^{-st}}{s^2 + a^2} (-s \cos at + a \sin at) \right]_0^{\infty} = \frac{s}{s^2 + a^2}, \text{ if } s > 0$$

$$(6) \quad L\{\sinh at\} = \int_0^{\infty} e^{-st} \sinh at dt = \int_0^{\infty} \frac{e^{at} - e^{-at}}{2} dt = \frac{1}{2} \left[\frac{e^{(s-a)t}}{s-a} - \frac{e^{-(s+a)t}}{s+a} \right]_0^{\infty} = \frac{a}{s^2 - a^2}, \text{ for } s > |a|$$

$$(7) \quad L\{\cosh at\} = \int_0^{\infty} e^{-st} \cosh at dt = \int_0^{\infty} \frac{e^{at} + e^{-at}}{2} dt = \frac{1}{2} \left[\frac{e^{(s-a)t}}{s-a} + \frac{e^{-(s+a)t}}{s+a} \right]_0^{\infty} = \frac{s}{s^2 - a^2}, \text{ for } s > |a|$$

PROPERTIES OF LAPLACE TRANSFORMS:

1. LINEARITY PROPERTY:

If a, b, c be any constants and f, g, h any functions of t , then

$$L\{af(t) + bg(t) - ch(t)\} = aL\{f(t)\} + bL\{g(t)\} - cL\{h(t)\}$$

By definition,

$$\begin{aligned} L.H.S. &= \int_0^{\infty} e^{-st} [af(t) + bg(t) - ch(t)] dt \\ &= a \int_0^{\infty} e^{-st} f(t) dt + b \int_0^{\infty} e^{-st} g(t) dt - c \int_0^{\infty} e^{-st} h(t) dt \\ &= aL\{f(t)\} + bL\{g(t)\} - cL\{h(t)\} \end{aligned}$$

II. FIRST SHIFTING PROPERTY:

If $L\{f(t)\} = \bar{f}(s)$, then

$$L\{e^{at}f(t)\} = \bar{f}(s-a)$$

By definition,

$$\begin{aligned} L\{e^{at}f(t)\} &= \int_0^{\infty} e^{-st} e^{at} f(t) dt = \int_0^{\infty} e^{-(s-a)t} f(t) dt \\ &= \int_0^{\infty} e^{-rt} f(t) dt, \text{ where } r = s-a \end{aligned}$$

$a = \bar{f}(r) = \bar{f}(s-a).$

APPLICATION OF FIRST SHIFTING PROPERTY:

- (1) $L\{e^{at}\} = \frac{1}{s-a}$
- (2) $L\{e^{at}t^n\} = \frac{n!}{(s-a)^{n+1}}$ when $n=1, 2, 3, \dots$
- (3) $L\{e^{at}\sin bt\} = \frac{b}{(s-a)^2 + b^2}$
- (4) $L\{e^{at}\cos bt\} = \frac{s-a}{(s-a)^2 + b^2}$
- (5) $L\{e^{at}\sinh bt\} = \frac{b}{(s-a)^2 - b^2}$
- (6) $L\{e^{at}\cosh bt\} = \frac{s-a}{(s-a)^2 - b^2}$ where in each case $a > b$.

III. CHANGE OF SCALE PROPERTY:

If $L\{f(t)\} = \bar{f}(s)$, then

$$L\{f(at)\} = \frac{1}{a} \bar{f}\left(\frac{s}{a}\right)$$

By definition,

$$L\{f(at)\} = \int_0^{\infty} e^{-st} f(at) dt$$

$$\begin{aligned} &= \int_0^{\infty} e^{-su/a} f(u) \frac{du}{a} \quad [\text{put } at = u, dt = \frac{du}{a}] \\ &= \frac{1}{a} \int_0^{\infty} e^{-su/a} f(u) du \\ &= \frac{1}{a} \bar{f}\left(\frac{s}{a}\right) \end{aligned}$$

Example–3:

Find the Laplace transform of $e^{2t}(3t^5 - \cos 4t)$.

Solution:

$$\begin{aligned} L\{e^{2t}(3t^5 - \cos 4t)\} &= 3L\{e^{2t}t^5\} - L\{e^{2t}\cos 4t\} \\ &= 3 \times \frac{5!}{(s-2)^6} - \frac{s-2}{(s-2)^2 + 4^2} = \frac{360}{(s-2)^6} - \frac{s-2}{s^2 - 4s + 20} \end{aligned}$$

Example–4:

Find the Laplace transform of $e^{-t}\sin^2 3t$

Solution:

We have

$$L\{\sin^2 3t\} = \frac{1}{2} L\{1 - \cos 6t\} = \frac{1}{2} \left[\frac{1}{s} - \frac{s}{s^2 + 36} \right] = \frac{18}{s(s+36)}$$

Example–5:

Find the Laplace transform of $e^{-3t}\sin 5t \sin 3t$.

Solution:

$$\text{We have, } L\{\sin 5t \sin 3t\} = \frac{1}{2} L\{\cos 2t - \cos 8t\}$$

$$= \frac{1}{2} \left[\frac{s}{s^2 + 2^2} - \frac{s}{s^2 + 8^2} \right] = \frac{30s}{(s+4)(s+64)}$$

By first shifting property, we get

Therefore, $L\hat{1} = \frac{\ddot{y}}{t} = \tan^{-1} \frac{a\ddot{a}\ddot{o}}{\zeta} = \frac{\ddot{a}\ddot{o}}{\ddot{s}\ddot{o}}$

LAPLACE TRANSFORMS OF DERIVATIVES:

(1) If $f(t)$ be continuous and $L\{f(t)\} = \bar{f}(s)$, then $L\{f'(t)\} = s\bar{f}(s) - f(0)$.

Proof: We have

$$L\{f'(t)\} = \int_0^{\infty} e^{-st} f'(t) dt$$

$$= \left[e^{-st} f(t) \right]_0^{\infty} - \int_0^{\infty} (-s) e^{-st} f(t) dt$$

Now assuming $f(t)$ be such that $\lim_{t \rightarrow \infty} e^{-st} f(t) = 0$, we have

$$L\{f'(t)\} = -f(0) + s \int_0^{\infty} e^{-st} f(t) dt$$

Thus, $L\{f'(t)\} = s\bar{f}(s) - f(0)$

(2) If $f(t)$ and its first $(n-1)$ derivatives be continuous, then

$$L\{f^{(n)}(t)\} = s^n \bar{f}(s) - s^{n-1} f(0) - s^{n-2} f'(0) - \dots - f^{(n-1)}(0)$$

Thus,

$$L\{f''(t)\} = s^2 \bar{f}(s) - s f(0) - f'(0)$$

$$L\{f'''(t)\} = s^3 \bar{f}(s) - s^2 f(0) - s f'(0) - f''(0)$$

$$L\{f^{(iv)}(t)\} = s^4 \bar{f}(s) - s^3 f(0) - s^2 f'(0) - s f''(0) - f'''(0) \text{ and so on.}$$

Laplace transforms of integrals:

$$\text{If } L\{f(t)\} = \bar{f}(s), \text{ then } L\left\{\int_0^t f(u) du\right\} = \frac{1}{s} \bar{f}(s)$$

Proof :

$$\text{Let } f(t) = \int_0^t f(u) du, \text{ then } f'(t) = f(t) \text{ and } f(0) = 0$$

$$L\{f'(t)\} = s \bar{f}(s) - f(0)$$

$$\text{or } L\{f(t)\} = \frac{\bar{f}(s)}{s}$$

$$\text{or } \bar{f}(s) = s \bar{f}(s)$$

$$\text{or } \bar{f}(s) = \frac{1}{s} \bar{f}(s)$$

$$\text{Hence, } L\left\{\int_0^t f(u) du\right\} = \frac{1}{s} \bar{f}(s).$$

Multiplication By t^n :

If $\mathcal{L}\{f(t)\} = f(s)$, then

$$\mathcal{L}\{t^n f(t)\} = (-1)^n \frac{d^n f(s)}{ds^n}, \text{ where } n=1, 2, 3, \dots$$

Division By t :

If $\mathcal{L}\{f(t)\} = f(s)$, then

$$\mathcal{L}\left\{\frac{f(t)}{t}\right\} = \int_s^\infty f(s) ds, \text{ provided the integral exists.}$$

Example–8:

Find the Laplace transforms of

(1) $t \sin at$

(2) $t \cos at$

Solution:

$$(1) \text{ We have, } \mathcal{L}\{t \sin at\} = \frac{d}{ds} \left(\frac{a}{s^2 + a^2} \right) = - \frac{2as}{(s^2 + a^2)^2}$$

$$\text{Hence } \mathcal{L}\{t \sin at\} = \frac{2as}{(s^2 + a^2)^2}$$

$$(2) \text{ We have, } \mathcal{L}\{t \cos at\} = \frac{d}{ds} \left(\frac{s}{s^2 + a^2} \right) = \frac{s^2 - a^2}{(s^2 + a^2)^2}$$

$$\text{Hence, } \mathcal{L}\{t \cos at\} = \frac{s^2 - a^2}{(s^2 + a^2)^2}$$

Example–9:

Find the Laplace transform of $t^2 \cos at$.

Solution:

$$\text{We have, } \mathcal{L}\{t \cos at\} = \frac{d}{ds} \left(\frac{s}{s^2 + a^2} \right) = \frac{s^2 - a^2}{(s^2 + a^2)^2}$$

$$\mathcal{L}\{t^2 \cos at\} = (-1)^2 \frac{d^2}{ds^2} \left(\frac{s}{s^2 + a^2} \right) = \frac{2s^2 - 2a^2}{(s^2 + a^2)^3}$$

$$\frac{=d\hat{e}a^2-s^2\hat{u}ds}{\hat{e}\hat{e}(s^2+a^2)^2\hat{u}\hat{u}}$$

Example–10:

Find the Laplace transforms of $\frac{(e^{-at} - e^{-bt})}{t}$.

Solution:

[illegible]

Example-11:

Find the inverse Laplace transform of $\frac{ae^{at} - \cos bt}{t}$.

Solution:

$$\text{We have, } L\hat{\mathbf{i}} \frac{\hat{\mathbf{i}} e^{at} \text{-cos} b t \hat{\mathbf{u}}}{t} \hat{\mathbf{y}} = \hat{\mathbf{o}} \hat{\mathbf{e}} \frac{1}{s + b \hat{\mathbf{u}}} - \frac{s}{s^2 + b^2} \hat{\mathbf{u}} ds$$

$$\begin{aligned} &= \hat{e}^{\dot{e}} \log(s-a) \cdot \frac{1}{2} \log(s_2+b_2) \hat{u}^{\infty} \\ &= \frac{1}{2} \hat{e}^2 \log(s-a) \cdot \log(s_2+b_2) \hat{u}^{\infty} \\ &= \frac{1}{2} \hat{e}^2 \hat{u}_s \\ &= \frac{1}{2} \log \zeta_s^2 \frac{1}{\hat{e} s + b} \frac{1}{\hat{u}} \\ &= \frac{1}{2} \hat{e} \log 1 - \log \zeta_s^2 \frac{1}{\hat{e} s + b} \frac{1}{\hat{u}} \\ &= \frac{1}{2} \log \zeta_s^2 (s-a)^2 \frac{1}{\hat{e} s + b} \frac{1}{\hat{u}} \\ &= \frac{1}{2} \log \hat{e}^2 \frac{1}{\hat{e} s + b} \frac{1}{\hat{u}} \end{aligned}$$

INVERSE LAPLACE TRANSFORMS:

We know that if $\{f(t)\} = f(s)$, then $L^{-1}\{f(s)\} = f(t)$

Let us now determine the inverse Laplace transforms of some given functions.

$$\begin{aligned}
(1) \quad & \begin{array}{c} {}_{-1}\hat{1}\ddot{u} \\ L\hat{1}-\hat{y}=1 \\ \hat{1}s\mathfrak{p} \end{array} \\
(2) \quad & \begin{array}{c} {}_{-1}\hat{1} \quad 1\ddot{u} \quad at \\ L\hat{1}-\hat{y}=e \\ \hat{1}s-a\mathfrak{p} \end{array} \\
(3) \quad & \begin{array}{c} {}_{-1}\hat{1}1\ddot{u} \quad t^{n-1} \\ L\hat{1}-\hat{y}=\frac{\quad}{n\hat{y}}, n=1, 2, 3, \dots \\ \hat{1}s\mathfrak{p} \quad (n-1)! \end{array} \\
(4) \quad & \begin{array}{c} {}_{-1}\hat{1} \quad 1 \quad \ddot{u} \quad et^{n-1} \\ L\hat{1}-\hat{y}=\frac{\quad}{n\hat{y}}, n=1, 2, 3, \dots \\ \hat{1}(s-a) \quad \mathfrak{p} \quad (n-1)! \end{array} \\
(5) \quad & \begin{array}{c} {}_{-1}\hat{1} \quad 1 \quad \ddot{u} \quad 1 \\ L\hat{1}-\hat{y}=\frac{\quad}{2} \hat{y} = -\sin at \\ \hat{1}s+a\mathfrak{p} \quad a \end{array} \\
(6) \quad & \begin{array}{c} {}_{-1}\hat{1} \quad 1 \quad \ddot{u} \\ L\hat{1}-\hat{y}=\frac{\quad}{2} \hat{y} = \cos at \\ \hat{1}s + a\mathfrak{p} \end{array} \\
(7) \quad & \begin{array}{c} {}_{-1}\hat{1} \quad 1 \quad \ddot{u}=1 \sin hat \\ L\hat{1}-\hat{y}=\frac{\quad}{2} \hat{y} = - \\ \hat{1}s-a\mathfrak{p} \quad a \end{array}
\end{aligned}$$

$$(8) \quad \mathcal{L}^{-1} \left\{ \frac{s}{s^2 - a^2} \right\} = \cosh at$$

$$(9) \quad \mathcal{L}^{-1} \left\{ \frac{1}{s^2 - a^2} \right\} = \frac{1}{a} \sinh at$$

$$\mathcal{L}^{-1} \left\{ \frac{s}{s^2 + b^2} \right\} = \cos bt$$

$$\mathcal{L}^{-1} \left\{ \frac{b}{s^2 + b^2} \right\} = \sin bt$$

$$(10) \quad \mathcal{L}^{-1} \left\{ \frac{1}{s^2 + a^2} \right\} = \frac{1}{a} \sin at$$

$$(11) \quad \mathcal{L}^{-1} \left\{ \frac{s}{(s+a)^2} \right\} = t e^{-at}$$

$$\mathcal{L}^{-1} \left\{ \frac{1}{(s+a)^2} \right\} = t e^{-at}$$

$$(12) \quad \mathcal{L}^{-1} \left\{ \frac{1}{s^2 + a^2} \right\} = \frac{1}{a} \sin at$$

$$\mathcal{L}^{-1} \left\{ \frac{s}{s^2 + a^2} \right\} = \cos at$$

INVERSE LAPLACE TRANSFORMS BY THE METHOD OF PARTIAL FRACTIONS:

We have seen that $\mathcal{L}\{f(t)\}$ in many cases, is a rational algebraic function of s . Hence to find the inverse Laplace transforms of $f(s)$, we first express the given function of s into partial fractions which will, then, be recognizable as one of the above mentioned standard forms.

Example–12:

Find the inverse Laplace transform of $\frac{s^2 + s + 2}{(s+1)^2(s-3)}$.

Solution:

Suppose that,

$$\frac{s^2 + s + 2}{(s+1)^2(s-3)} = \frac{A}{s+1} + \frac{B}{(s+1)^2} + \frac{C}{s-3} \quad \dots(1)$$

Multiplying both sides of (1) by $(s+1)^2(s-3)$, we get

$$s^2 + s + 2 = A(s+1)(s-3) + B(s-3) + C(s+1)^2 \quad \dots(2)$$

Putting $s = -1$

$$2 = -4B \quad B = -\frac{1}{4}$$

Putting $s = 3$

$$14 = 16C \quad C = \frac{7}{8}$$

$$14 = 16C \quad C = \frac{7}{8}$$

Equating coefficients of s^2 , we get

$$1 = A + C$$

$$1 = A + \frac{7}{8} \quad A = 1 - \frac{7}{8} = \frac{1}{8}$$

Engineering Mathematics–III

Putting the values of A, B, C in (1) we get

$$\frac{s^2 + s + 2}{(s+1)^2 (s-3)^2} = \frac{1}{(s+1)^2} + \frac{1}{s-3} + \frac{7}{(s-3)^2}$$

$$\mathcal{L}^{-1} \left\{ \frac{s^2 + s + 2}{(s+1)^2 (s-3)^2} \right\} = \mathcal{L}^{-1} \left\{ \frac{1}{(s+1)^2} \right\} + \mathcal{L}^{-1} \left\{ \frac{1}{s-3} \right\} + \mathcal{L}^{-1} \left\{ \frac{7}{(s-3)^2} \right\}$$

$$= \frac{1}{8} e^{-t} - \frac{1}{8} t e^{-t} + e^{3t} + \frac{7}{8} t e^{3t}$$

Example–13:

Find the inverse Laplace transform of $\frac{s}{(s-2)(s^2+9)}$

Solution:

Suppose that,

$$\frac{s}{(s-2)(s^2+9)} = \frac{A}{s-2} + \frac{Bs+C}{s^2+9} \quad \dots(1)$$

Multiplying both sides by $(s-2)(s^2+9)$, we get

$$s = A(s^2+9) + (Bs+C)(s-2) \quad \dots(2)$$

$$\text{Putting } s=2, 2=13A \Rightarrow A=\frac{2}{13}$$

$$\text{Putting } s=0, 0=9A-2C \Rightarrow C=\frac{9}{13}$$

Equating coefficients of s^2 , we get

$$0 = A + B \Rightarrow B = -\frac{2}{13}$$

Putting the values of A, B, C in (1), we get

$$\frac{s}{(s-2)(s^2+9)} = \frac{2}{13} \frac{1}{s-2} - \frac{2}{13} \frac{s}{s^2+9} + \frac{9}{13} \frac{1}{s^2+9}$$

$$\mathcal{L}^{-1} \left\{ \frac{s}{(s-2)(s^2+9)} \right\} = \frac{2}{13} \mathcal{L}^{-1} \left\{ \frac{1}{s-2} \right\} - \frac{2}{13} \mathcal{L}^{-1} \left\{ \frac{s}{s^2+9} \right\} + \frac{9}{13} \mathcal{L}^{-1} \left\{ \frac{1}{s^2+9} \right\}$$

$$= \frac{2}{13} e^{2t} - \frac{2}{13} \cos 3t + \frac{3}{13} \sin 3t$$

OTHER METHODS OF FINDING INVERSE LAPLACE TRANSFORMS:**(I) SHIFTING PROPERTY:**

If $\mathcal{L}^{-1}\{f(s)\} = f(t)$, then

$$\mathcal{L}^{-1}\{f(s-a)\} = e^{at}f(t) = e^{at}\mathcal{L}^{-1}\{f(s)\}$$

(II) If $\mathcal{L}^{-1}\{f(s)\} = f(t)$ and $f(0) = 0$, then $\mathcal{L}^{-1}\{sf(s)\} = \frac{d}{dt}f(t)$.

$$d^n$$

In general, $\mathcal{L}^{-1}\{s^n f(s)\} = \frac{d^n}{dt^n}f(t)$, Provided f

$$(0) = f(0) = \dots = f^{(n-1)}(0) = 0.$$

$$\mathcal{L}^{-1}\left\{s \frac{d}{ds} f(s)\right\} = -f(t), \text{ then } \mathcal{L}^{-1}\left\{s^2 \frac{d^2}{ds^2} f(s)\right\} = f(t)$$

(IV) If $\mathcal{L}^{-1}\{f(s)\} = f(t)$, then $f(t) = \mathcal{L}^{-1}\left\{\frac{d}{ds} f(s)\right\}$.**(V) If $f(t) = \mathcal{L}^{-1}\{f(s)\}$, then $\mathcal{L}\left\{\frac{d}{dt}f(t)\right\} = -sf(s)$.**

This formula is useful in finding $f(t)$ when $f(s)$ is given.

Example–14:

Find the inverse Laplace transform of $\frac{2s}{s^2+4}$.

$$\frac{2s}{s^2+4}$$

Solution:

$$\text{Let } \mathcal{L}^{-1}\left\{\frac{2s}{s^2+4}\right\} = f(t)$$

$$\mathcal{L}\{f(t)\} = \frac{2s}{s^2+4} = f(s)$$

Then by formula IV we get,

$$\mathcal{L}\{t f(t)\} = -\frac{d}{ds} f(s) = -\frac{d}{ds} \left(\frac{2s}{s^2+4}\right)$$

$$= -2 \frac{(s^2+4) - s(2s)}{(s^2+4)^2}$$

$$= \frac{2}{s^2+4}$$

Engineering Mathematics–III

$$\mathcal{L}^{-1}\left\{\frac{1}{s^2+4}\right\} = \sin 2t$$

$$\mathcal{L}^{-1}\left\{\frac{\sin 2t}{t}\right\}$$

$$\mathcal{L}^{-1}\left\{\frac{1}{s^2+4}\right\} = \sin 2t$$

Example–16:

Find the inverse Laplace transform of $\log \frac{s}{s+1}$.

Solution:

$$\mathcal{L}^{-1}\left\{\log \frac{s}{s+1}\right\} = f(t)$$

$$\mathcal{L}\{f(t)\} = \log \frac{s}{s+1}$$

Then by formula IV we get,

$$\mathcal{L}\{f(t)\} = \int_0^\infty \log \frac{s}{s+1} ds = \int_0^\infty (\log s - \log(s+1)) ds$$

$$= -\frac{1}{s} - \frac{1}{s+1} = -\frac{1}{s} - \frac{1}{s+1}$$

$$\mathcal{L}^{-1}\left\{-\frac{1}{s} - \frac{1}{s+1}\right\} = -1 - e^{-t}$$

$$\mathcal{L}^{-1}\left\{-\frac{1}{s} - \frac{1}{s+1}\right\} = -1 - e^{-t}$$

$$\mathcal{L}^{-1}\left\{\log \frac{s}{s+1}\right\} = \frac{e^{-t} - 1}{t}$$

Example–17:

Find the inverse Laplace transform of $\frac{1}{s^2(s^2+a^2)}$.

Solution:

We have

$$\mathcal{L}^{-1}\left\{\frac{1}{s^2(s^2+a^2)}\right\} = \frac{1}{a^2} \sin at = f(t).$$

Then by Formula III we get,

$$L^{-1}\left\{\frac{1}{s^2(s^2+a^2)}\right\} = \frac{1}{a} \int_0^t \sin at \, dt = \frac{1}{a} \left[-\frac{\cos at}{a} \right]_0^t = \frac{1}{a^2} (1 - \cos at)$$

Thus we have,

$$\begin{aligned} L^{-1}\left\{\frac{1}{s^2(s^2+a^2)}\right\} &= \frac{1}{a^2} \int_0^t (1 - \cos at) \, dt \\ &= \frac{1}{a^2} \left[t - \frac{\sin at}{a} \right]_0^t \\ &= \frac{1}{a^2} (at - \sin at) \end{aligned}$$

Example – 18:

Find the inverse Laplace transform of $\frac{1}{s^2s+5}$.

Solution:

$$\text{We have, } L^{-1}\left\{\frac{1}{s^2s+5}\right\} = f(t).$$

Then by Formula III, we get

$$\begin{aligned} L^{-1}\left\{\frac{1}{s^2(s+5)}\right\} &= \frac{1}{5} \int_0^t (1 - e^{-5t}) \, dt \\ &= \frac{1}{5} \left[t - \frac{1 - e^{-5t}}{-5} \right]_0^t \\ &= \frac{1}{5} \left[t + \frac{1 - e^{-5t}}{5} \right] \end{aligned}$$

Thus we have,

$$\begin{aligned} L^{-1}\left\{\frac{1}{s^2(s+5)}\right\} &= \frac{1}{5} \int_0^t (1 - e^{-5t}) \, dt \\ &= \frac{1}{5} \left[t + \frac{1 - e^{-5t}}{5} \right] \\ &= \frac{1}{5} t + \frac{1 - e^{-5t}}{25} \end{aligned}$$

Engineering Mathematics–III

$$= \frac{1}{5} e^{-5t} + \frac{1}{5} e^{-5t} - \frac{1}{5} e^{-5t}$$

$$= \frac{1}{5} e^{-5t} + \frac{1}{5} e^{-5t} - \frac{1}{5} e^{-5t}$$

Example–19:

Find the inverse Laplace transform of $\frac{s^2}{(s^2 + a^2)^2}$

Solution:

We have,

$$L^{-1} \left\{ \frac{s^2}{s^2 + a^2} \right\} = \frac{1}{2a} \frac{d}{dt} (t \sin at) = f(t)$$

$$L^{-1} \left\{ \frac{s^2}{s^2 + a^2} \right\} = \frac{1}{2a} \frac{d}{dt} (t \sin at)$$

Since $f(0) = 0$, we get from Formula II that,

$$L^{-1} \left\{ \frac{s^2}{s^2 + a^2} \right\} = \frac{1}{2a} \frac{d}{dt} (t \sin at) = \frac{1}{2a} (\sin at + at \cos at)$$

Example–20:

Find the inverse Laplace transform of $\frac{s+3}{(s^2+6s+13)^2}$.

Solution:

$$\text{We have, } \frac{s+3}{(s^2+6s+13)^2} = \frac{s+3}{(s^2+6s+13)^2} = \frac{s+3}{(s^2+6s+13)^2}$$

Then by formula I we get

$$L^{-1} \left\{ \frac{s+3}{(s^2+6s+13)^2} \right\} = L^{-1} \left\{ \frac{s+3}{(s^2+6s+13)^2} \right\} = L^{-1} \left\{ \frac{s+3}{(s^2+6s+13)^2} \right\}$$

$$= e^{-3t} \cdot t \sin 2t$$

Example–21:

Find the inverse Laplace transform of $\frac{s}{(s^2 + a^2)^2}$.

Solution:

$$\text{Let, } f(t) = L^{-1} \left\{ \frac{s}{(s^2 + a^2)^2} \right\}$$

Then by formula V we get

$$L^{-1} \left\{ \frac{s}{(s^2 + a^2)^2} \right\} = \frac{1}{2} L^{-1} \left\{ \frac{2s}{(s^2 + a^2)^2} \right\}$$

$$= \frac{1}{2} L^{-1} \left\{ \frac{1}{s^2 + a^2} - \frac{1}{(s^2 + a^2)^2} \right\}$$

$$= \frac{1}{2} \left[L^{-1} \left\{ \frac{1}{s^2 + a^2} \right\} - L^{-1} \left\{ \frac{1}{(s^2 + a^2)^2} \right\} \right]$$

$$= \frac{1}{2} \left[\frac{1}{a} \sin at - \frac{1}{a^2} t \sin at \right]$$

$$\text{Hence } f(t) = \frac{1}{2a} \sin at - \frac{1}{2a^2} t \sin at$$

LAPLACE TRANSFORM METHOD TO SOLVE LINEAR DIFFERENTIAL EQUATIONS WITH CONSTANT COEFFICIENTS ASSOCIATED WITH INITIAL CONDITIONS :

Linear differential equations with constant coefficients associated with initial conditions can be easily solved by Laplace transform method.

Working Procedure:

Step - 1 : Take the Laplace transform of both sides of the differential equation and then put the given initial conditions.

Step-2: Transpose the terms with minus sign to the right.

Step-3: Divide by the coefficient of $\bar{y}$, getting $\bar{y}$ as a known function of s .

Step-5: Resolve this function of s into partial fractions.

Step–5 : Take the inverse Laplace transform of both sides. This gives y as a function of t which is the desired solution satisfying the given conditions.

Example–22:

Solve the following equation by transform method; $y'' - 3y' + 2y = e^{3t}$, when $y(0) = 1$ and $y'(0) = 0$.

Solution:

We have, $y'' - 3y' + 2y = e^{3t}$... (1)

Taking Laplace transform of both sides of (1), we get

$$L\{y''\} = 3L\{y'\} + 2L\{y\} = L\{e^{3t}\}$$

$$\Rightarrow [s^2 \bar{y} - sy(0) - y'(0)] - 3[s\bar{y} - y(0)] + 2\bar{y} = \frac{1}{s-3}$$

Putting $y(0) = 1$ and $y'(0) = 0$, we get

$$s^2 \bar{y} - s - 3s\bar{y} + 3 + 2\bar{y} = \frac{1}{s-3}$$

$$\Rightarrow \bar{y}(s^2 - 3s + 2) = \frac{1 + (s-3)^2}{s-3}$$

$$\Rightarrow \bar{y} = \frac{s^2 - 6s + 10}{(s-3)(s^2 - 3s + 2)}$$

$$\Rightarrow \bar{y} = \frac{s^2 - 6s + 10}{(s-3)(s-1)(s-2)} \quad \dots (2)$$

$$\text{Let, } \frac{s^2 - 6s + 10}{(s-3)(s-1)(s-2)} = \frac{A}{s-3} + \frac{B}{s-1} + \frac{C}{s-2}$$

Multiplying both sides of (3) by $(s-3)(s-1)(s-2)$, we get

$$s^2 - 6s + 10 = A(s-1)(s-2) + B(s-3)(s-2) + C(s-3)(s-1) \quad \dots (4)$$

$$\begin{aligned} \text{Putting } s=1, B &= \frac{5}{2} \\ \text{Putting } s=2, C &= -2 \end{aligned}$$

$$\text{Putting } s=3, A = \frac{1}{2}$$

Substituting the values of A, B, C in (3), we get

$$\bar{y} = \frac{1}{2} \frac{1}{s-3} + \frac{5}{2} \frac{1}{s-1} - 2 \frac{1}{s-2}$$

Taking inverse Laplace transform of both sides, we get

$$y = \frac{1}{2} e^{3t} + \frac{5}{2} e^t - 2e^{2t}$$

This is the required solution.

Example–23:

Solve the following equation by transform method;

$$(D^2 + \omega^2)y = \cos \omega t, t > 0, \text{ given that } y=0 \text{ and } Dy=0 \text{ at } t=0$$

Solution:

We have

$$(D^2 + \omega^2)y = \cos \omega t$$

i.e., $y'' + \omega^2 y = \cos \omega t$, given $y(0) = y'(0) = 0$ Taking Laplace transform of both sides of (1), we get

$$L\{y''\} + \omega^2 L\{y\} = L\{\cos \omega t\}$$

$$p \quad s^2 y - sy(0) - y'(0) + \omega^2 y = \frac{s}{s^2 + \omega^2}$$

Putting $y(0) = 0$ and $y'(0) = 0$, we get

$$y(s^2 + \omega^2) = \frac{s}{s^2 + \omega^2}$$

$$p \quad y = \frac{s}{s^2 + \omega^2} \quad \dots(2)$$

Taking inverse Laplace transform, we get

$$L^{-1}\{y\} L^{-1}\left\{\frac{s}{s^2 + \omega^2}\right\}$$

$$y = \frac{1}{\omega} \sin \omega t$$

This is the required solution.

Assignment

1. Find the Laplace transforms of the following:

$$\begin{array}{ll} \text{(a)} \quad \frac{R_t}{T_0} \quad \frac{U}{W} & \text{(b)} \quad \frac{R_t}{T_0} \quad \frac{U}{W} \end{array}$$

2. Find the Laplace Transform of $f(t)$ in each of the following:

$$\begin{array}{ll} \text{(a)} \quad f(t) = \begin{cases} S \sin 2t, & \text{when } 0 < t \leq \pi \\ T_0, & \text{when } t > \pi \end{cases} & \text{(b)} \quad f(t) = \begin{cases} S e^{-t}, & \text{when } 0 \leq t \leq 2 \\ T_0, & \text{when } t > 2 \end{cases} \end{array}$$

3. Obtain the inverse Laplace transforms of the following functions

$$\begin{array}{ll} \text{(a)} \quad \frac{2s^2 - 6s + 5}{s^3 - 6s^2 + 11s - 6} & \text{(b)} \quad \frac{a(s^2 - 2a^2)}{s^4 + 4a^4} \end{array}$$

$$\begin{array}{ll} \text{(c)} \quad \frac{s}{s^2 + 6s + 13} & \text{(d)} \quad \log \frac{F(1+s)I}{G(s)H(s)K} \end{array}$$

rrr

CHAPTER–4**FOURIER SERIES****Periodic Functions:**

If the value of each ordinate $f(t)$ repeat itself at equal interval in the abscissa, then $f(t)$ is said to be a periodic function.


If $f(t) = f(t+T) = f(t+2T) = \dots$, then

T is called period of the function $f(t)$.

For example

$$\sin x = \sin(x+2\pi) = \sin(x+4\pi) = \dots$$

So $\sin x$ is called a periodic function of period 2π

**Fourier Series:**

A series of sines and cosines of an angle and its multiple of the form

$$\frac{a_0}{2} + a_1 \cos x + \frac{a_2}{2} \cos 2x + \dots + a_n \cos nx + \dots$$

$$+ b_1 \sin x + b_2 \sin 2x + \dots + b_n \sin nx + \dots$$

$$= \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx$$

is called a Fourier series, there a_0 , a_n & b_n are called Fourier constants **Useful Integrals**

The following integrals are useful in Fourier series:

$$1. \int_0^{2\pi} \sin nx \, dx = 0$$

$$2. \int_0^{2\pi} \cos nx \, dx = 0$$

$$3. \quad \frac{d}{dx} \int_a^{a+2\pi} \sin^2 nx dx = p$$

$$4. \quad \frac{d}{dx} \int_a^{a+2\pi} \cos^2 nx dx = p$$

$$5. \quad \frac{d}{dx} \int_a^{a+2\pi} \sin nx \cdot \sin mx dx = 0$$

$$6. \quad \frac{d}{dx} \int_a^{a+2\pi} \cos nx \cdot \cos mx dx = 0$$

$$7. \quad \frac{d}{dx} \int_a^{a+2\pi} \sin nx \cdot \cos mx dx = 0$$

$$8. \quad \frac{d}{dx} \int_a^{a+2\pi} \sin nx \cdot \sin mx dx = 0$$

$$9. \quad \frac{d}{dx} uv = u \frac{dv}{dx} + v \frac{du}{dx} + u^2 v^3 - \dots$$

$$\text{where } v_1 = \frac{du}{dx}, v_2 = \frac{d^2u}{dx^2}, v_3 = \frac{d^3u}{dx^3} \dots$$

$$10. \quad \sin p = 0 \text{ and } \cos p = (-1)^n \text{ where } n \in \mathbb{I}$$

Let $f(x)$ be represented in the interval $(a, a+2p)$ by Fourier series

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx \quad \dots(1)$$

To find a_0 :

Integrate both sides of equation (1) from $x=a$ to $x=a+2p$. Then

$$\begin{aligned} \int_a^{a+2p} f(x) dx &= \int_a^{a+2p} \left[\frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx \right] dx \\ &= \frac{1}{2} a_0 (a+2p-a) + 0 + 0 = a_0 p \end{aligned}$$

$$\text{Hence } a_0 = \frac{1}{p} \int_a^{a+2p} f(x) dx$$

To find a_n : Multiply $\cos nx$ on both sides of equation (1) and integrate from $x=a$ to $x=a+2p$, Then

$$\begin{aligned} \int_a^{a+2p} f(x) \cos nx dx &= \int_a^{a+2p} \left[\frac{a_0}{2} \cos nx + \sum_{n=1}^{\infty} a_n \cos nx \cos nx + \sum_{n=1}^{\infty} b_n \sin nx \cos nx \right] dx \\ &= \frac{a_0}{2} \int_a^{a+2p} \cos nx dx + \sum_{n=1}^{\infty} a_n \int_a^{a+2p} \cos^2 nx dx + \sum_{n=1}^{\infty} b_n \int_a^{a+2p} \sin nx \cos nx dx \\ &= 0 + p a_n + 0 \end{aligned}$$

$$a_n = \frac{1}{p} \int_a^{a+2p} f(x) \cos nx dx$$

To find b_n : Multiply $\sin nx$ on both sides of equation (1) and integrate from $x=a$ to $x=a+2p$, then

$$\begin{aligned} \int_a^{a+2p} f(x) \sin nx dx &= \int_a^{a+2p} \left[\frac{a_0}{2} \sin nx + \sum_{n=1}^{\infty} a_n \cos nx \sin nx + \sum_{n=1}^{\infty} b_n \sin nx \sin nx \right] dx \\ &= 0 + 0 + p b_n \end{aligned}$$

Engineering Mathematics–III

$$\text{Hence } b_n = -\frac{1}{\pi} \int_{\alpha}^{\alpha+2\pi} f(x) \sin nx \, dx$$

Making $\alpha=0$, the interval becomes $0 < x < p$ and the formula (1) reduces to

$$\begin{aligned} a_0 &= \frac{1}{p} \int_0^{2\pi} f(x) \, dx \\ a_n &= \frac{1}{p} \int_0^{2\pi} f(x) \cos nx \, dx \quad \dots \text{(ii)} \\ b_n &= \frac{1}{p} \int_0^{2\pi} f(x) \sin nx \, dx \end{aligned}$$

Putting $\alpha=-p$, The interval becomes $-p < x < p$, the formula (I) reduces to

$$\begin{aligned} a_0 &= \frac{1}{p} \int_{-p}^p f(x) \, dx \\ a_n &= \frac{1}{p} \int_{-p}^p f(x) \cos nx \, dx \quad \dots \text{(iii)} \\ b_n &= \frac{1}{p} \int_{-p}^p f(x) \sin nx \, dx \end{aligned}$$

Euler's Formula:

The Fourier series for the function $f(x)$ in the interval $-p < x < p+2\pi$ is given by

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx$$

$$\text{where } a_0 = \frac{1}{p} \int_{\alpha}^{\alpha+2\pi} f(x) \, dx$$

$$a_n = \frac{1}{p} \int_{\alpha}^{\alpha+2\pi} f(x) \cos nx \, dx$$

$$b_n = \frac{1}{p} \int_{\alpha}^{\alpha+2\pi} f(x) \sin nx \, dx$$

The value of a_0 , a_n & b_n are known.

Euler's formula.

Example–1:

Given that $f(x) = x + x^2$ for $-p < x < p$, find the Fourier expansion of $f(x)$. Hence that

$$\frac{\pi^2}{6} = 1 + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots$$

Solution:

$$\text{Let } x = \frac{a_0}{2} + a_1 \cos x + a_2 \cos 2x + \dots + b_1 \sin x + b_2 \sin 2x + \dots \quad \dots(1)$$

$$a_0 = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) dx = \frac{1}{\pi} \int_{-\pi}^{\pi} (x + x^2) dx = \frac{1}{\pi} \left[\frac{x^2}{2} + \frac{x^3}{3} \right]_{-\pi}^{\pi} = \frac{1}{\pi} \left[\frac{\pi^2}{2} + \frac{\pi^3}{3} - \left(\frac{(-\pi)^2}{2} + \frac{(-\pi)^3}{3} \right) \right] = \frac{1}{\pi} \left[\frac{\pi^2}{2} + \frac{\pi^3}{3} - \frac{\pi^2}{2} + \frac{\pi^3}{3} \right] = \frac{2\pi^2}{3}$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx = \frac{1}{\pi} \int_{-\pi}^{\pi} (x + x^2) \cos nx dx$$

$$= \frac{1}{\pi} \left[\frac{x \sin nx}{n} - \frac{(-\cos nx)}{n^2} + \frac{x^2 \sin nx}{2n} - \frac{2x \cos nx}{n^2} + \frac{2 \sin nx}{n^3} \right]_{-\pi}^{\pi}$$

$$= \frac{1}{\pi} \left[\frac{x \sin nx}{n} - \frac{\cos nx}{n^2} + \frac{x^2 \sin nx}{2n} - \frac{2x \cos nx}{n^2} + \frac{2 \sin nx}{n^3} \right]_{-\pi}^{\pi}$$

$$= \frac{1}{\pi} \left[\frac{\pi \sin n\pi}{n} - \frac{\cos n\pi}{n^2} + \frac{\pi^2 \sin n\pi}{2n} - \frac{2\pi \cos n\pi}{n^2} + \frac{2 \sin n\pi}{n^3} \right] - \left[\frac{(-\pi) \sin n\pi}{n} - \frac{\cos n\pi}{n^2} + \frac{(-\pi)^2 \sin n\pi}{2n} - \frac{2(-\pi) \cos n\pi}{n^2} + \frac{2 \sin n\pi}{n^3} \right]$$

$$= \frac{1}{\pi} \left[\frac{\pi \sin n\pi}{n} - \frac{\cos n\pi}{n^2} + \frac{\pi^2 \sin n\pi}{2n} - \frac{2\pi \cos n\pi}{n^2} + \frac{2 \sin n\pi}{n^3} + \frac{\pi \sin n\pi}{n} - \frac{\cos n\pi}{n^2} - \frac{\pi^2 \sin n\pi}{2n} + \frac{2\pi \cos n\pi}{n^2} - \frac{2 \sin n\pi}{n^3} \right]$$

$$= \frac{1}{\pi} \left[\frac{2\pi \sin n\pi}{n} - \frac{2 \cos n\pi}{n^2} \right] = \frac{2 \sin n\pi}{n} - \frac{2 \cos n\pi}{n^2}$$

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx$$

$$= \frac{1}{\pi} \left[\frac{x \cos nx}{n} - \frac{(-\sin nx)}{n^2} + \frac{x^2 \cos nx}{2n} - \frac{2x \sin nx}{n^2} + \frac{2 \cos nx}{n^3} \right]_{-\pi}^{\pi}$$

$$= \frac{1}{\pi} \left[\frac{x \cos nx}{n} - \frac{\sin nx}{n^2} + \frac{x^2 \cos nx}{2n} - \frac{2x \sin nx}{n^2} + \frac{2 \cos nx}{n^3} \right]_{-\pi}^{\pi}$$

$$= \frac{1}{\pi} \left[\frac{\pi \cos n\pi}{n} - \frac{\sin n\pi}{n^2} + \frac{\pi^2 \cos n\pi}{2n} - \frac{2\pi \sin n\pi}{n^2} + \frac{2 \cos n\pi}{n^3} \right] - \left[\frac{(-\pi) \cos n\pi}{n} - \frac{\sin n\pi}{n^2} + \frac{(-\pi)^2 \cos n\pi}{2n} - \frac{2(-\pi) \sin n\pi}{n^2} + \frac{2 \cos n\pi}{n^3} \right]$$

$$= \frac{1}{\pi} \left[\frac{\pi \cos n\pi}{n} - \frac{\sin n\pi}{n^2} + \frac{\pi^2 \cos n\pi}{2n} - \frac{2\pi \sin n\pi}{n^2} + \frac{2 \cos n\pi}{n^3} + \frac{\pi \cos n\pi}{n} - \frac{\sin n\pi}{n^2} - \frac{\pi^2 \cos n\pi}{2n} + \frac{2\pi \sin n\pi}{n^2} - \frac{2 \cos n\pi}{n^3} \right]$$

$$= \frac{1}{\pi} \left[\frac{2\pi \cos n\pi}{n} - \frac{2 \sin n\pi}{n^2} \right] = \frac{2 \cos n\pi}{n} - \frac{2 \sin n\pi}{n^2}$$

Substituting the values of a_0, a_n & b_n in equation (1)

$$x + x^2 = \frac{2\pi^2}{3} + 4 \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} \cos 2nx + \sum_{n=1}^{\infty} \frac{2 \sin n\pi}{n} - \frac{2 \cos n\pi}{n^2} \sin 3x + \dots \quad \dots(2)$$

Put $x = \pi$ in equation (2)

EngineeringMathematics–III

$$p+p \rightarrow p^2 + 4^1 e^- + \frac{1}{2} + \frac{1}{2} + \frac{1}{4} + \dots \rightarrow \dots (3)$$

$$\text{Putx} = -\text{pinequation} \quad (2)$$

$$-p+p^2 = \frac{p^2}{3} + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots + \frac{1}{n} \quad (4)$$

Addingequation(3)&(4)

$$\begin{array}{ccccccc}
2p^2 & \acute{e} & 1 & 1 & 1 & \grave{u} \\
2p = \frac{\quad}{3} + 8\acute{e}1+ & \frac{\quad}{2+} & \frac{\quad}{2+2} & \frac{\quad}{\quad} & \frac{\quad}{\quad} & + \dots \dots \acute{u} \\
& \grave{e} & 2 & 3 & 4 & \hat{u} \\
4p^2 & \acute{e} & 1 & 1 & 1 & \grave{u} \\
\frac{\quad}{\quad} = 8\acute{e}1+2+2+ & \frac{\quad}{\quad} & \frac{\quad}{\quad} & \frac{\quad}{2+} & \dots \dots \dots \acute{u} \\
& 3 & \grave{e} & 2 & 3 & 4 \\
p^2 & \underline{1} & \underline{1} & \underline{1} & \hat{u} & \frac{1}{\infty} \\
\frac{\quad}{\quad} & & & = 1+2+2+2+ & \dots \dots = \hat{a}_2 & \\
6 & 2 & 3 & 4 & & \overset{X}{\underset{n=1}{\hat{a}_2}}
\end{array}$$

Dirchelet'sCondition:

$$\text{Any function } f(x) \text{ can be developed as a fourier series } \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx$$

where a_0, a_n, b_n are constants provided.

- (i) $f(x)$ is periodic, single valued and finite
- (ii) $f(x)$ has a finite no. of discontinuities in any one period
- (iii) $f(x)$ has at most a finite no. of maxima and minima.

Discontinuous Functions: At a point of discontinuity, Fourier series gives the value of $f(x)$ as the arithmetic mean of left and right limits.

At a point of discontinuity, $x=c$

$$f(x) = \frac{1}{2} [f(c-0) + f(c+0)]$$

Example-2:

Find the fourier series expansion for

$$\begin{array}{l} \text{Deducethat} \\ \frac{1}{1} + \frac{1}{3} + \frac{1}{5} + \dots = \frac{p^2}{8} \end{array}$$

Solution:

$$\text{Let } f(x) = \frac{a_0}{Z} + \sum_{n=1}^{\infty} \dot{a}_n \cos nx + \sum_{n=1}^{\infty} \dot{b}_n \sin nx \dots (1)$$

$$\begin{aligned}
 \text{then } a_0 &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx \\
 &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \{x\} dx = \frac{1}{2\pi} \left[\frac{x^2}{2} \right]_{-\pi}^{\pi} = \frac{1}{2\pi} \left(\frac{\pi^2}{2} - \frac{(-\pi)^2}{2} \right) = 0 \\
 a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx \\
 &= \frac{1}{\pi} \int_{-\pi}^{\pi} \{x\} \cos nx dx \\
 &= \frac{1}{\pi} \left[\frac{x \sin nx}{n} - \int \sin nx dx \right]_{-\pi}^{\pi} \\
 &= \frac{1}{\pi} \left[\frac{x \sin nx}{n} + \frac{\cos nx}{n} \right]_{-\pi}^{\pi} \\
 &= \frac{1}{\pi} \left[\frac{\pi \sin n\pi}{n} + \frac{\cos n\pi}{n} - \left(\frac{-\pi \sin n\pi}{n} + \frac{\cos n\pi}{n} \right) \right] \\
 &= \frac{1}{\pi} \left[\frac{\cos n\pi}{n} - \frac{\cos n\pi}{n} \right] = 0
 \end{aligned}$$

$$\begin{aligned}
 a_1 &= -\frac{2}{\pi}, a_2 = 0, a_3 = -\frac{2}{\pi}, a_4 = 0 \\
 b_1 &= \frac{1}{\pi}, b_2 = -\frac{1}{\pi}, b_3 = \frac{1}{\pi}, b_4 = -\frac{1}{\pi} \\
 \text{Substituting the values of } a_n \text{ and } b_n \text{ in equation (1), we get} \\
 f(x) &= \frac{1}{2} - \frac{\cos 3x}{4} + \frac{\cos 5x}{8} - \frac{\cos 7x}{16} + \dots \\
 &+ \frac{\sin 2x}{3} - \frac{\sin 4x}{6} + \frac{\sin 6x}{9} - \dots \quad \text{..... (ii)}
 \end{aligned}$$

Putting $x=0$ in equation (ii)

$$\begin{aligned}
 f(0) &= \frac{1}{2} - \frac{\cos 0}{4} + \frac{\cos 0}{8} - \frac{\cos 0}{16} + \dots \\
 &+ \frac{\sin 0}{3} - \frac{\sin 0}{6} + \frac{\sin 0}{9} - \dots \quad \text{..... (iii)}
 \end{aligned}$$

Now $f(x)$ is discontinuous at $x=0$ But

$$f(0-0) = f(0+0) = 0$$

$$f(0) = \frac{1}{2} [f(0-0) + f(0+0)] = \frac{1}{2} [0 + 0] = 0$$

Engineering Mathematics–III

From equation (iii)

$$-\frac{p}{24p^6} = -\frac{p}{1^2} + \frac{1}{2} + \frac{1}{5^2} + \dots$$


or $\frac{p}{8} = -\frac{1}{1} + \frac{1}{3} + \frac{1}{5} + \dots$

Even Function: A function $f(x)$ is said to be even (or symmetric) function if $f(-x) = f(x)$

Ex. (i) $x^2, x^4, x^6, \dots$ even powers of x

(ii) $\cos x, \sec x$ etc.

The graph of such a function is symmetrical with respect to y-axis. Here y-axis is a mirror for the reflection of the curve




The area under such a curve from $-p$ to p is double the area from 0 to p .

$$\int_{-p}^p f(x) dx = 2 \int_0^p f(x) dx$$

Odd function: A function $f(x)$ is called odd (or skew symmetric) function if $f(-x) = -f(x)$

Ex. (i) $x^3, x^5, x^7, \dots$ odd powers of x

(ii) $\sin x, \csc x, \tan x$ etc



Here the area under the curve from $-p$ to p is zero

$$\int_{-p}^p f(x) dx = 0$$

Expansion of an Even Function:

$$a_0 = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) dx = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) dx$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx = \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx dx$$

As $f(x)$ and $\cos nx$ both are even, the product of $f(x) \cdot \cos nx$ is also even.

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx = 0$$

As $\sin nx$ is an odd function. The product of an even function with odd function is odd, therefore we need not calculate b_n .

The series of an even function contains cosine terms only.

Expansion of an odd Function:

$$a_0 = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) dx = 0$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx = 0 \quad \text{Q } (f(x) \cdot \cos nx \text{ is odd function})$$

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx = \frac{2}{\pi} \int_0^{\pi} f(x) \sin nx dx$$

(Q $f(x) \cdot \sin nx$ is even function)

The series of an odd function contains sine terms only.

Example–3:

Obtain a Fourier expansion of $f(x) = x^3$ in $-\pi < x < \pi$

Solution:

$f(x) = x^3$ is an odd function.

$$a_0 = 0 \text{ and } a_n = 0$$

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} x^3 \sin nx dx = \frac{2}{\pi} \int_0^{\pi} x^3 \sin nx dx$$

$$= \frac{2}{\pi} \left[-x^3 \cos nx + 3x^2 \sin nx - 6x \cos nx + 6 \sin nx \right]_0^{\pi}$$

$$= \frac{2}{\pi} \left[-\pi^3 \cos n\pi + 3\pi^2 \sin n\pi - 6\pi \cos n\pi + 6 \sin n\pi \right]$$

$$= \frac{2}{\pi} \left[-\pi^3 (-1)^n + 3\pi^2 (0) - 6\pi (-1)^n + 6(0) \right]$$

$$= \frac{2}{\pi} \left[\pi^3 (-1)^{n+1} - 6\pi (-1)^n \right]$$

$$= 2(-1)^{n+1} \pi^2 + 12(-1)^n$$

$$f(x) = \frac{1}{2} \left[f(x) + f(x) \right] = \frac{1}{2} \left[\sum_{n=1}^{\infty} b_n \sin nx + \sum_{n=1}^{\infty} b_n \sin nx \right] = \sum_{n=1}^{\infty} b_n \sin nx$$

Half Range Series:

To obtain a Fourier expansion of a function $f(x)$ for the range $(0, p)$ which is half the period of the fourier series. As it is immaterial whatever the function may be outside the range $0 < x < p$, we extend the function to cover the range $-p < x < p$. So that the new function may be even or odd. The fourier expansion of such function of half the period consists sine or cosine term only.

Sine Series:

If it is required to expand $f(x)$ as a sine series in $0 < x < p$ we extend the function to the range $-p < x < p$, so that it will be an odd function.

The desired half-range sine series is given by

$$f(x) = \sum_{n=1}^{\infty} b_n \sin nx$$

$$\text{where } b_n = \frac{2}{p} \int_0^p f(x) \sin nx dx$$

Cosine Series:

If it is required to expand $f(x)$ as a cosine series in $0 < x < p$, We extend the function to the range $-p < x < p$, so that it will be an even function.

The desired half-range cosine series is given by

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos nx$$

$$\text{where } a_0 = \frac{2}{p} \int_0^p f(x) dx$$

$$a_n = \frac{2}{p} \int_0^p f(x) \cos nx dx$$

Example–4:

Find the half-range sine series for the function $f(x) = e^{ax}$ for $0 < x < p$

Solution:

$$f(x) = \sum_{n=1}^{\infty} b_n \sin nx$$

$$\text{where } b_n = \frac{2}{p} \int_0^p f(x) \sin nx dx = \frac{2}{p} \int_0^p e^{ax} \sin nx dx$$

$$\begin{aligned}
 &= -\frac{e^{ax}}{2} \left[(a \sin nx - n \cos nx) \right]_{\hat{u}_0}^{\hat{u}} \\
 &= -\frac{e^{ax}}{2} \left[(a \sin np - n \cos np) \right] + \frac{n}{a^2 + n^2} \hat{u} \\
 &= \frac{2n}{a^2 + n^2} e^{ax} \cos nx + \frac{1}{a^2 + n^2} e^{ax} \sin nx \\
 &= \frac{2n}{(a^2 + n^2)^2} e^{ax} \cos nx + \frac{1}{(a^2 + n^2)^2} e^{ax} \sin nx \\
 &= \frac{2n(1 + e^{a\pi})}{(a^2 + 1)^2} \cos nx + \frac{2.2(1 - e^{a\pi})}{(a^2 + 2^2)^2} \sin nx \\
 &= \frac{2n(1 + e^{a\pi})}{(a^2 + 1)^2} \cos nx + \frac{2(1 - e^{a\pi})}{(a^2 + 2)^2} \sin nx
 \end{aligned}$$

Assignment

- Find a Fourier series to represent $f(x) = p - x, 0 < x < 2p$
- Find a Fourier series to represent the function $f(x) = e^x$, for $-p < x < p$

- Find the Fourier series of the function $f(x) = \begin{cases} 0, & \text{for } -p < x < -\frac{p}{2} \\ 1, & \text{for } -\frac{p}{2} < x < \frac{p}{2} \\ 0, & \text{for } \frac{p}{2} < x < p \end{cases}$
- Represent the following function by a Fourier sine series $f(t) = \begin{cases} 1, & \text{for } 0 < t < \frac{p}{2} \\ 0, & \text{for } \frac{p}{2} < t < p \end{cases}$
- Find the Fourier cosine series for the function $f(x) = \begin{cases} 1, & \text{for } -p < x < -\frac{p}{2} \\ 0, & \text{for } -\frac{p}{2} < x < \frac{p}{2} \\ 1, & \text{for } \frac{p}{2} < x < p \end{cases}$

CHAPTER–5**FINITE DIFFERENCE AND INTERPOLATION****Finite Difference:**

Suppose we are given the following values of $y = f(x)$ for a set of values x :

x	x_0	x_1	$x_2, \dots$	x_n
y	y_0	y_1	$y_2, \dots$	y_n

The interpolation is the technique of estimating the value of a function for any intermediate value of the independent variable. While the process of computing the value of the function outside the given range is called extrapolation.

Suppose that the function $y = f(x)$ is tabulated for the equally spaced values $x = x_0, x_0 + h, x_0 + 2h, \dots, x_0 + nh$ giving $y = y_0, y_1, \dots, y_n$. To determine the values of $f(x)$ for some intermediate values of x , the following two types of difference are found useful.

Forward difference - The differences

$$Dy_0 = y_1 - y_0$$

$$Dy_1 = y_2 - y_1$$

$$Dy_{n-1} = y_n - y_{n-1}$$

Similarly

$$D^2y_0 = Dy_1 - Dy_0$$

$$D^3y_1 = D^2y_2 - D^2y_1$$

$$D^n y_0 = D^{n-1} y_1 - D^{n-1} y_0$$

Forward difference table

Value of x	Value of y	1st diff.	2nd diff.	3rd diff.	4th diff.	5th diff.
x_0	y_0	Dy_0				
$x+h$ ₀	y ₁	Dy ₁	D^2y ₀			
$x+2h$ ₀	y ₂	Dy ₂	D^2y ₁	D^3y ₀		
$x+3h$ ₀	y ₃	Dy ₃	D^2y ₂	D^3y ₁	D^4y ₀	
$x+4h$ ₀	y ₄	Dy ₄	D^2y ₃	D^3y ₂	D^4y ₁	D^5y ₀
x_0+5h	y_5					

Backward difference

$$\tilde{N}y_1 = y_1 - y_0$$

$$\tilde{N}y_1 = y_1 - y_0$$

:

:

:

$$\tilde{N}^n y_n = \tilde{N}^{n-1} y_n - \tilde{N}^{n-1} y_{n-1}$$

Backward difference table

Value of x	Value of y	1st diff.	2nd diff.	3rd diff.	4th diff.	5th diff.
x_0	y_0	Dy_0				
$x_0 + h$	y_1	Dy_1	D^2y_0			
$x_0 + 2h$	y_2	Dy_2	D^2y_1	D^3y_0		
$x_0 + 3h$	y_3	Dy_3	D^2y_2	D^3y_1	D^4y_0	
$x_0 + 4h$	y_4	Dy_4	D^2y_3	D^3y_2	D^4y_1	D^5y_0
$x_0 + 5h$	y_5					

Differences of a polynomial

We know that the expression of the form $f(x) = a_0x^n + a_1x^{n-1} + \dots + a_{n-1}x + a_n$ where a 's are constant ($a_0 \neq 0$) and n is a positive integer is called a polynomial in x of degree n .

Theorem:

The 1st difference is a polynomial of degree $n-1$, the 2nd difference is of degree $n-2$, and the n th difference is constant. While higher differences are equal to zero.

The converse of the theorem is also true which states that if n th difference of a function tabulated at equally spaced intervals are constant, the function is a polynomial of degree n .

Example-1:

Form the successive forward differences of ax^3 , the interval being h .

Solution:

$$\text{Here } y = f(x) = ax^3$$

We know that

$$Dy_0 = y_1 - y_0 = f(x+h) - f(x)$$

$$\backslash D(ax^3) = a(x+h)^3 - ax^3$$

$$= a(x^3 + 3x^2h + 3xh^2 + h^3) - ax^3$$

$$= a(3x^2h + 3xh^2 + h^3)$$

$$\text{Again, } D^2y_0 = Dy_1 - Dy_0$$

$$\backslash D^2(ax^3) = a\{3(x+h)^2h + 3(x+h)h^2 + h^3\} - a(3x^2h + 3xh^2 + h^3)$$

$$= a\{3x^2h + 6xh^2 + 3h^3 + 3xh^2 + 3h^3 + h^3 - 3x^2h - 3xh^2 - h^3\} = a\{6xh^2 + 6h^3\}$$

Engineering Mathematics–III

$$D^3 y_0 = D^2 y_1 = D^2 y_0$$

$$\backslash D^3(ax^3) = a\{6(x+h)h^2 + 6h^3\} - a\{6xh^2 + 6h^3\}$$

$$= a\{6xh^2 + 6h^3 + 6h^3 - 6xh^2 - 6h^3\} = 6ah^3 = \text{Constant}$$

$$\backslash D^4(ax^3) = 6ah^3 - 6ah^3 = 0$$

Here it shows that the the third differences of a polynomial of third degree is constant & the higher difference & are zero.

Factorial Notation

A product of the form $x(x-1)(x-2)\dots\dots\dots(x-r+1)$ is devoted by $[x]^r$ and is called a factorial.

In particular

$$[x] = x, [x]^2 = x(x-1)$$

$$[x]^3 = x(x-1)(x-2)$$

$$[x]^n = x(x-1)(x-2)\dots\dots\dots(x-n+1)$$

which is called a factorial polynomial or function.

The factorial notation is of special utility in the theory of first differences. It helps in finding the successive differences of a polynomial directly by simply rule of differentiation. The result of differentiating $[x]^r$ is similar to that of differential x^r .

Example–2:

Estimate the missing term in the following table:

x	0	1	2	3	4
$f(x)$	1	3	9	8	1

Solution:

Let the missing term be y_3 . The following is the difference table.

x	y	D	D^2	D_3	D^4
0	1				
1	3	2	4	$y_3 - 19$	
2	9	6	$y_3 - 15$		$124 - 4y_3$
3	y_3	$y_3 - 9$	$81 - 2y_3 + 9$	$105 - 3y_3$	
4	81	$81 - y_3$			

As only four entries y_0, y_1, y_2, y_4 are given, the function y can be represented by a third degree polynomial, here 4th order difference becomes zero, i.e.,

$$124 - 4y_3 = 0$$

$$\Rightarrow y_3 = 31$$

Hence the missing term is 31.

Example–3:

Estimate the missing term in the following table:

x	0	1	2	3	4	5	6
y	5	11	22	40	--	140	--

Solution:

Let the missing term be y_4 & y_6 . The following is the difference table.

x:	y:	D	D ₂	D ₃	D ₄	D ₅
0	5					
1	11	6	5			
2	22	11	7	2	$y_4 - 67$	
3	40	18	$y_4 - 40$	$y_4 - 58$	$303 - 4y_4$	$370 - 5y_4$
4	y_4	$y_4 - 40$	$180 - 3y_4$	$238 - 3y_4$	$y_6 + 6y_4 -$	$y_6 + 10y_4 - 1001$
5	140	$140 - y_4$	$y_6 + y_4 -$	$y_6 + 3y_4 - 460$	698	
6	y_6	$y_6 - 140$	280			

As only four entries y_0, y_1, y_2, y_4, y_5 are given, the function y can be represented by a 4th degree polynomial & hence 5th difference becomes zero, i.e.,

$$370 - 5y_4 = 0 \quad \text{and} \quad y_6 + 10y_4 - 1001 = 0$$

Solving these, we get

$$y_4 = 74 \quad \text{and} \quad y_6 = 261$$

Newton's Forward interpolation formula for equal intervals

Let the function $y = f(x)$ take the values $y_0, y_1, y_2, \dots$ corresponding to the values

0 1 2

$x_0, x_1 + h, x_0 + 2h$ of x .

$$f(x_0 + nh) = y_0 + nDy_0 + \frac{n(n-1)}{2!} D^2y_0 + \frac{n(n-1)(n-2)}{3!} D^3y_0 + \dots$$

Obs. This formula is used for interpolating the values of y near the beginning of a set of tabulated values and extrapolating values of y a little backward (i.e. to the left) of y_0 .

Newton's backward interpolation formula for equal intervals

Let the function $y = f(x)$ take the values $y_0, y_1, y_2, \dots$ corresponding to the values

0 1 2

$x_0, x_1 + h, x_0 + 2h$ of x .

$$f(x_n + nh) = y_n + n\tilde{N}y_n + \frac{n(n+1)}{2!} \tilde{N}^2y_n + \frac{n(n+1)(n+2)}{3!} \tilde{N}^3y_n + \dots$$

Obs. This formula is used for interpolating the values of y near the beginning of a set of tabulated values and extrapolating values of y a little backward (i.e. to the right) of y_n .

Example–4:

The table gives the distances in nautical miles of the visible horizon for the given heights in feet above the earth's surface.

x=height	100	150	200	250	300	350	400
y=distance	10.63	13.03	15.04	16.81	18.42	19.90	21.27

Find the values of y when (i) $x=218$ ft.

Solution:

The difference table is as under:

x	y	D	D ₂	D ₃	D ₄
100	10.63				
		2.40			
150	13.03		–0.39		
		2.01		0.15	
200	15.04		–0.24		–0.07
		1.77		0.08	
250	16.81		–0.16		–0.05
		1.61		0.03	
300	18.42		–0.13		–0.01
		1.48		0.02	
350	19.90		–0.11		
		1.37			
400	21.27				

- (i) If we take $x_0=200$, then $y_0=15.04$, $Dy_0=1.77$, $D^2y_0=–0.16$, $D^3y_0=0.03$ etc.

$$\text{Since } x=218 \text{ and } h=50, \therefore n = \frac{x-x_0}{h} = \frac{18}{50} = 0.36$$

Using Newton's forward interpolation formula, we get

$$y_{218} = y_0 + n\Delta y_0 + \frac{n(n-1)}{1.2} \Delta^2 y_0 + \frac{n(n-1)(n-2)}{1.2.3} \Delta^3 y_0 + \dots$$

$$f(218) = 15.04 + 0.36(1.77) + \frac{0.36(-0.64)}{26} + \frac{0.36(-0.64)(-1.64)}{(-0.16) + \dots} + (0.03) + \dots$$

- (ii) Since $x=410$ is near the end of the table, we use Newton's backward interpolation formula.

$$\text{Taking } x_n=400, n = \frac{x-x_n}{h} = \frac{10}{50} = 0.2$$

Using the line of backward differences

$$Y_n = 21.27, \tilde{N}^2 y_n = -0.11, \tilde{N}^3 y_n = 0.02 \text{ etc.}$$

Newton's backward formula gives

$$y_{410} = y_{400} + n\tilde{N}y_{200} + \frac{n(n+1)}{2}\tilde{N}^2y_{400} + \frac{n(n+1)(n+2)}{1.2.3}\tilde{N}^3y_{400} + \dots$$

$$= 21.27 + 0.2(1.37) + \frac{0.2(1.2)}{2}(-0.11) + \dots = 21.53 \text{ nautical miles.}$$

Example–5:

Find the number of men getting wages between Rs. 10 and 15 from the following data:

Wages in Rs.	0–10	10–20	20–30	30–40
Frequency	9	30	35	42

Solution:

First we prepare the cumulative frequency table, as follows:

Wages less than (x)	10	20	30	40
No. of men (y)	9	39	74	116

Now the difference table is

x	y	D	D ²	D ³
10	9			
		30		
20	39		5	
		35		2
30	74		7	
		42		
40	116			

We shall find y_{15} i.e. number of men getting wages less than 15.

Taking $x_0 = 10$, $x = 15$, we have

$$n = \frac{x - x_0}{h} = \frac{15 - 10}{10} = 0.5$$

Using Newton's forward interpolation formula, we get

$$y_{15} = y_{10} + nDy_{10} + \frac{n(n-1)}{2!}D^2y_{10} + \frac{n(n-1)(n-2)}{3!}D^3y_{10}$$

$$= 9 + (0.5)'30 + \frac{(0.5)(0.5-1)}{2}'5 + \frac{(0.5)(0.5-1)(0.5-2)}{6}'2$$

$$= 9 + 15 - 0.625 + 0.125 = 23.5 = 24 \text{ approx.}$$

Number of men getting wages between Rs. 10 and 15 = 24 – 10 = 14 approx.

Example–6:

Find the cubic polynomial which takes the following values:

x	0	1	2	3
$f(x)$	1	2	1	10

Solution:

The difference table is

x	$f(x)$	$Df(x)$	$D^2f(x)$	$D^3f(x)$
0	1			
		1		
1	2		-2	
		-1		12
2	1		10	
		9		
3	10			

We take $x_0 = 0$ and $p = \frac{x - x_0}{h} = x$

Using Newton's forward interpolation formula, we get

$$\begin{aligned}
 f(x) &= f(x_0) + \frac{x(x-1)}{1 \cdot 2} Df(0) + \frac{x(x-1)(x-2)}{1 \cdot 2 \cdot 3} D^2f(0) + \frac{x(x-1)(x-2)(x-3)}{1 \cdot 2 \cdot 3 \cdot 4} D^3f(0) \\
 &= 1 + \frac{x(x-1)}{2} (-2) + \frac{x(x-1)(x-2)}{6} (12) \\
 &= 2x^3 + 7x^2 + 6x + 1, \text{ which is the required polynomial.}
 \end{aligned}$$

To compute $f(4)$, we take $x_n = 3, x = 4$ so that $p = \frac{x - x_n}{h} = 1$

Using Newton's backward interpolation formula, we get

$$\begin{aligned}
 f(4) &= f(x_n) + \frac{n}{1} \tilde{N} f(x_n) + \frac{n(n+1)}{1 \cdot 2} \tilde{N}^2 f(x_n) + \frac{n(n+1)(n+2)}{1 \cdot 2 \cdot 3} \tilde{N}^3 f(x_n) \\
 &= 10 + 9 + 10 + 12 + 41
 \end{aligned}$$

which is the same value as that obtained by substituting $x = 4$ in the cubic polynomial above.

Obs. The above example shows that if a tabulated function is a polynomial, then interpolation and extrapolation give the same values.

Lagrange's Interpolation formula for unequal intervals:

$$f(x) = \frac{(x-x_1)(x-x_2)\dots(x-x_n)}{(x-x_1)(x_0-x_2)\dots(x_0-x_n)}$$

$$y = \frac{(x-x_1)(x-x_2)\dots\dots\dots(x-x_n)}{(x_1-x_0)(x_1-x_2)\dots\dots\dots(x_1-x_n)}$$

$$y_1 + \dots\dots\dots - \frac{(x-x_0)(x-x_1)\dots\dots\dots(x-x_{n-1})}{(x_n-x_0)(x_n-x_2)\dots\dots\dots(x_n-x_{n-1})}$$

Lagrange's Method for unequally spaced values of x:

$$x = \frac{(y-y_1)(y-y_2)\dots\dots\dots}{(y_0-y_1)(y_0-y_2)\dots\dots\dots} + \frac{(y-y_n)}{(y_0-y_n)} + \frac{(y-y_0)(y-y_2)\dots\dots\dots}{(y_1-y_0)(y_1-y_2)\dots\dots\dots} + \dots\dots\dots + \frac{(y-y_0)(y-y_1)\dots\dots\dots}{(y_n-y_0)(y_n-y_1)\dots\dots\dots} + \frac{(y-y_{n-1})}{(y_n-y_{n-1})}$$

Example–7:

Use lagrange's interpolation formula to find the value of y when x = 10, if the following values of x & y are given.

Solution:

Here	x ₀ =5	x ₁ =6	x ₂ =9	x ₃ =11
and	y ₀ =12	y ₁ =13	y ₂ =14	y ₃ =16

Putting x=10 and substituting the above value in Lagrange's formula, we get:

$$f(x) = \frac{(x-x_1)(x-x_2)(x-x_3)}{(x_0-x_1)(x_0-x_2)(x_0-x_3)} y_0 + \frac{(x-x_0)(x-x_2)(x-x_3)}{(x_1-x_0)(x_1-x_2)(x_1-x_3)} y_1 + \frac{(x-x_0)(x-x_1)(x-x_3)}{(x_2-x_0)(x_2-x_1)(x_2-x_3)} y_2 + \frac{(x-x_0)(x-x_1)(x-x_2)}{(x_3-x_0)(x_3-x_1)(x_3-x_2)} y_3$$

$$\begin{aligned} f(10) &= \frac{(10-6)(10-9)(10-11) \times 12}{5(5-9)(5-11)(5-6)} + \frac{(10-5)(10-9)(10-11) \times 13}{6(6-5)(6-9)(6-11)} \\ &\quad + \frac{(10-5)(10-6)(10-11) \times 14}{5(9-6)(9-11)(10-5)} + \frac{(10-5)(10-6)(10-9) \times 16}{5(9-6)(9-11)(10-5)} \\ &= \frac{4 \times 1 \times (-1)}{(-1) \times (-4) \times (-6)} \times 12 + \frac{5 \times 1 \times (-1)}{1 \times (-3) \times (-5)} \times 13 \\ &\quad + \frac{5 \times 4 \times (-1) \times 14}{4 \times 3 \times (-2) \times 6 \times 5} + \frac{5 \times 4 \times 1 \times 16}{2 \times 24 \times 15} - \frac{48}{-24} + \frac{-65}{60} + 320 \\ &= 2 - 4.33 + 11.66 + 5.33 = 14.66 \end{aligned}$$

Example–8:

Apply Lagrange's method to find the value of x when $f(x)=15$ from the given data.

x	5	6	9	11
$f(x)$	12	13	14	16

Solution:

Here

$$x_0=5, \quad x_1=6, \quad x_2=9, \quad x_3=11$$

$$y_0=12, \quad y_1=13, \quad y_2=14, \quad y_3=16$$

Taking $y=15$ and using the above results in Lagrange's inverse interpolation formula.

$$\begin{aligned}
 x=f(x) &= \frac{(y-y_1)(y-y_2)(y-y_3)}{(y-y_0)(y-y_1)(y-y_2)}x_0 + \frac{(y-y_0)(y-y_2)(y-y_3)}{(y-y_1)(y-y_2)(y-y_3)}x_1 \\
 &+ \frac{(y-y_0)(y-y_1)(y-y_3)}{(y_2-y_0)(y_2-y_1)(y_2-y_3)}x_2 + \frac{(y-y_0)(y-y_1)(y-y_2)}{(y_3-y_0)(y_3-y_1)(y_3-y_2)}x_3 \\
 &= \frac{(15-13)(15-14)(15-16)}{(13-13)(12-14)(12-16)} \times 5 + \frac{(15-12)(15-14)(15-16)}{(13-12)(13-14)(13-16)} \times 6 \\
 &\quad + \frac{(15-12)(15-13)(15-16) \times 9 + (15-12)(15-13)(15-14) \times 11}{(12)(14-13)(14-16)(16-12)(16-13)(16-16)}(14) \\
 &= \frac{2 \times 1 \times (-1)}{(-1) \times (-2) \times (-4)} \times 5 + \frac{3 \times 1 \times (-1)}{1 \times (-1) \times (-3)} \times 6 + \frac{3 \times 2 \times (-1)}{2 \times 1 \times (-2)} \times 9 + \frac{3 \times 2 \times 1}{4 \times 3 \times 2} \times 11 \\
 &= \frac{5}{-6} + \frac{27}{-6} + \frac{11}{-6} = 1.25 - 6 + 13.5 + 2.75 = 17.5 - 6 = 11.5
 \end{aligned}$$

Assignment

1. Find a cubic polynomial which takes the following values

x	0	1	2	3
$f(x)$	1	2	1	10

2. Given the values

x	5	7	11	13	17
y	15	0	3	2	14

Evaluate y_9 using Lagrange's formula.

3. Given $\sin 45^\circ = 0.7071$, $\sin 50^\circ = 0.7660$, $\sin 55^\circ = 0.8192$ and $\sin 60^\circ = 0.8660$. Find $\sin 52^\circ$ using Newton's forward interpolation formula.

CHAPTER–6**NUMERICAL SOLUTION OF EQUATION**

1. An expression of the form

$$f(x) = a_0x^n + a_1x^{n-1} + \dots + a_{n-1}x + a_n$$

where $a_0, a_1, a_2, \dots, a_n \neq 0$ are constants and n is a positive integer is called a polynomial in x of degree n .

2. The polynomial $f(x) = 0$

For example (1) $2x^2 + x^2 - 13x + 6 = 0$

$$(2) x^3 - 4x + 9 = 0$$

are called algebraic equations.

3. Transcendental equation-

If $f(x)$ is a function other than algebraic function such as trigonometric, logarithmic, exponential etc. then $f(x)$ is called transcendental function.

4. Root of an equation-

The value of x which satisfies $f(x) = 0$ is called the root of the equation.

Geometrically a root of the equation $f(x) = 0$ & $y = 0$ is the value of x where the graph meets the y -axis.

5. Solution of an equation-

The process of finding a root of an equation is known as the solution of an equation.

6. Different methods to solve the equations.

(a) Analytical method

(b) Graphical method

(c) Numerical method

7. Limitation of analytical method

This method produces very exact and accurate results. But it fails in many cases such as it fails to find roots of transcendental equations.

8. Limitation of graphical method-

This method is simple but these methods produce results to a low degree of accuracy.

9. Advantages of Numerical method-

This method is often of a repetitive nature. These consist in repeated execution of the same process. Where each step the result of the preceding step is used. This is known as an iteration process and is repeated till the result is obtained to a desired degree of accuracy.


The following are some Numerical methods to find root of algebraic and transcendental equation –

- (1) Bisection method
- (2) Newton–Raphson method

Bisection method:

This method consists of locating a root of the equation $f(x)=0$ between a and b . If $f(x)$ is continuous between a and b , and $f(a)$ and $f(b)$ are of opposite signs then there is a root between a and b . From the graph $f(a)$ is negative and $f(b)$ is positive then there is a root lies between a and b . The first approximation to the root is

$$x_1 = \frac{a+b}{2}$$



if $f(x) = 0$, then x_1 is the root of equation $f(x) = 0$. Otherwise the root lies between a and x_1 or x_1 and b according to $f(x_1)$ is positive or negative. Then we bisect the interval and continue the process until the root is found to desired accuracy.


In the fig. 1 $f(x_1)$ is +ve, so the root lies between a and x_1 . Then the 2nd approximation to the root is $x_2 = \frac{1}{2} (a + x_1)$. If $f(x)$ is -ve, the root lies between x_1 and x . So the third approximation to the root is $x_3 = \frac{1}{2} (x_1 + x_2)$ and so on. 2

Example–1:

(a) Find a root of the equation

$x^3 - 4x - 9 = 0$ using the bisection method correct to three decimal places.

Solution:



Let $f(x) = x^3 - 4x - 9$
 $f(2) = (2)^3 - 4(2) - 9 = -9$ (-ve)
 $f(3) = (3)^3 - 4(3) - 9 = 6$ (+ve)
 a root lies between 2 and 3.

First approximation to the root is

$$x_1 = \frac{1}{2}(2+3) = 2.5$$

$$f(x_1) = (2.5)^3 - 4(2.5) - 9 = -3.375 \text{ (-ve)}$$

the root lies between x_1 and 3. The second approximation to the root is

$$x_2 = \frac{1}{2}(x_1 + 3) = \frac{1}{2}(2.5 + 3) = 2.75$$

$$f(x_2) = (2.75)^3 - 4(2.75) - 9 = 0.7969 \text{ (+ve)}$$

the root lies between 2.5 and 2.75

$$x_3 = \frac{1}{2}(2.5 + 2.75) = 2.625$$

$$f(2.625) = (2.625)^3 - 4(2.625) - 9 = -1.4121 \text{ (-ve)}$$

the root lies between 2.625 and 2.75.

$$x_4 = \frac{1}{2}(2.625 + 2.75) = 2.6875$$

Repeating this process, the successive approximations are

$$x_5 = 2.71875, x_6 = 2.70313, x_7 = 2.71094, x_8 = 2.70703, x_9 = 2.70508,$$

$$x_{10} = 2.70605, x_{11} = 2.70654, x_{12} = 2.70642. \text{ Hence the root is } 2.7064.$$

Example–2:

Find the root of the equation $x \log_{10} x = 1.2$ which lies between 2 and 3, using bisection method taking 2 stages.

Solution:

$$\begin{aligned} \text{Let } f(x) &= x \log_{10} x - 1.2 = 2 \times \log_{10} 2 - 1.2 \\ &= 2 \times .3010 - 1.2 = -0.5979 \text{ (-ve)} \end{aligned}$$

$$f(3) = 3 \times \log_{10} 3 - 1.2 = 0.2314 \text{ (+ve)}$$

the root lies between 2 and 3

$$x_1 = \frac{1}{2}(2+3) = 2.5$$

$$f(2.5) = 2.5(\log_{10} 2.5) - 1.2 = -0.205 \text{ (-ve)}$$

the root lies between 2.5 and 3

$$x_2 = \frac{1}{2}(2.5 + 3) = 2.75$$

Hence the root is 2.75.

Example–3:

By using the bisection method, find an approximate root of the equation $\sin x = \frac{1}{x}$, that lies between $x = 1$ and $x = 1.5$ (measured in radians). Carry out computations upto the 7th stage.

Solution.

Let $f(x) = x \sin x - 1$. We know that $V = 57.3^\circ$

Since $f(1) = 1 \times \sin(1) - 1 = \sin(57.3^\circ) - 1 = -0.15849$

and $f(1.5) = 1.5 \times \sin(1.5) - 1 = 1.5 \times \sin(85.95^\circ) - 1 = 0.49625$;

a root lies between 1 and 1.5.

1

\first approximation to the root is $x_1 = 2(1 + 1.5) = 1.25$.

Then $f(x_1) = (1.25) \sin(1.25) - 1 = 1.25 \sin(71.625^\circ) - 1 = 0.18627$ and $f(1) < 0$.

\a root lies between 1 and $x_1 = 1.25$.

1

Thus the second approximation to the root is $x_2 = 2(1 + 1.25) = 1.125$.

Then $f(x_2) = 1.125 \sin(1.125) - 1 = 1.125 \sin(64.46^\circ) - 1 = 0.01509$ and $f(1) < 0$.

\a root lies between 1 and $x_2 = 1.125$.

1

Thus the third approximation to the root is $x_3 = 2(1 + 1.125) = 1.0625$

Then $f(x_3) = 1.0625 \sin(1.0625) - 1 = 1.0625 \sin(60.88^\circ) - 1 = -0.0718 < 0$

and $f(x_2) > 0$, i.e. now the root lies between $x_4 = 1.0625$ and $x_2 = 1.125$.

1

\ fourth approximation to the root is $x_4 = 2(1.0625 + 1.125) = 1.09375$ Then

$f(x_4) = -0.02836 < 0$ and $f(x_2) > 0$,

i.e., the root lies between $x_4 = 1.09375$ and $x_2 = 1.125$.

1

\ fifth approximation to the root is $x_5 = 2(1.09375 + 1.125) = 1.10937$

Then $f(x_5) = -0.00664 < 0$ and $f(x_2) > 0$.

\the root lies between $x_5 = 1.10937$ and $x_2 = 1.125$.

Thus the sixth approximation to the root is

1


$x_6 = 2(1.10937 + 1.125) = 1.11719$

Then $f(x_6) = 0.00421 > 0$. But $f(x_5) < 0$.

\the root lies between $x_5 = 1.10937$ and $x_6 = 1.11719$.

1

Thus the seventh approximation to the root is $x_7 = 2(1.10937 + 1.11719) = 1.11328$ Hence the desired approximation to the root is 1.11328.



In this method, instead of taking two initial rough approximations to the root $x=a$ as in the previous two methods, a single rough approximation x_0 to the root is taken. Then we use the following formula, known as Newton-Raphson formula or Newton iteration formula, to get the successive approximations.

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \quad \dots (1)$$

Putting $n=0, 1, 2, \dots$ etc. in the above formula (1), we get the first, second, third approximations as follows.

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$$

$$x_2 = x_1 - \frac{f(x_1)}{f'(x_1)}$$

$$x_3 = x_2 - \frac{f(x_2)}{f'(x_2)}$$

This method is useful in cases of large values of $f'(x)$ i.e., when the graph of $f(x)$ while crossing the x -axis is nearly vertical.

The process of finding successive approximations to the root (i.e., x_1, x_2, x_3 etc.) may be continued till the root is found to desired degree of accuracy.

Example–4:

Find by Newton's method, a root of the equation $x^3 - 3x + 1 = 0$ correct to 3 decimal places.

Solution:

$$\text{Let } f(x) = x^3 - 3x + 1$$

$$f(1) = 1 - 3 + 1 = -1$$

$$f(2) = 2^3 - 3 \cdot 2 + 1 = 8 - 6 + 1 = 3$$

∴ The root of $f(x)$ lies between 1 & 2

Engineering Mathematics–III

Let $x_0 = 1.5$, Also $f(x) = 3x^2 - 3$

Newton's formula gives

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} = x_n - \frac{(x_n^3 - 3x_n + 1)}{2x_n - 3}$$

$$= \frac{3x_n^3 - 3x_n - 3x_n^3 + 3x_n - 1}{3x_n^2 - 3} = \frac{2x_n^3 - 1}{3x_n^2 - 3} \quad \dots(1)$$

Putting $n=0$ in (i), the first approximation x_1 is given by

$$x_1 = \frac{2x_0^3 - 1}{3x_0^2 - 3} = \frac{2 \times (1.5)^3 - 1}{3 \times (1.5)^2 - 3} = \frac{2 \times 3.375 - 1}{3 \times 2.25 - 3} = \frac{5.75}{3.75} = 1.533$$

Putting $n=1$ in (i), the second approximation x_2 is given by

$$x_2 = \frac{2x_1^3 - 1}{3x_1^2 - 3} = \frac{2 \times (1.533)^3 - 1}{3 \times (1.533)^2 - 3} = \frac{2 \times 3.6026 - 1}{3 \times 2.35 - 3} = \frac{6.2052}{4.05} = 1.532$$

Example–5:

Find the Newton's method, the real root of the equation $3x = \cos x + 1$

Solution:

Let $f(x) = 3x - \cos x - 1$

$$f(0) = -2 = -ve, f(1) = 3 - 0.5403 - 1 = 1.4597 = +ve$$

So a root of $f(x) = 0$ lies between 0 and 1. It is nearer to 1. Let us take $x_0 = 0.6$. Also $f(x) = 3 + \sin x$

Newton's iteration formula gives

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} = x_n - \frac{3x_n - \cos x_n - 1}{3 + \sin x_n} = \frac{x_n \sin x_n + \cos x_n}{3 + \sin x_n} + 1 \quad \dots(i)$$

Putting $n=0$, the first approximation x_1 is given by

$$x_1 = \frac{x_0 \sin x_0 + \cos x_0 + 1}{3 + \sin x_0} = \frac{(0.6) \sin(0.6) + \cos(0.6) + 1}{3 + \sin(0.6)}$$

$$= \frac{0.6 \times 0.5729 + 0.82533 + 1}{3 + 0.5729} = 0.6071$$

Putting $n=1$ in (i), the second approximation is

$$x_2 = \frac{x_1 \sin x_1 + \cos x_1 + 1}{3 + \sin x_1} = \frac{0.6071 \sin(0.6071) + \cos(0.6071) + 1}{3 + \sin(0.6071)}$$

$$= \frac{0.6071 \times 0.57049 + 0.8213 + 1}{3 + 0.57049} = 0.6071$$

Clearly, $x_2 = x_1$.

Hence the desired root is 0.6071 correct to four decimal places.

Assignment

1. Find a root of the following equations, using the bisection method correct to three decimal places.
 - (a) $x^3 - x - 11 = 0$
 - (b) $x^4 - x - 10 = 0$
2. Find by Newton-Raphson method, a root of the following equations correct to 3 decimal places.
 - (a) $x^3 - 3x + 1 = 0$
 - (b) $3x^3 - 9x^2 + 8 = 0$
3. Using Newton-Raphson method to evaluate the following
 - (a) $\frac{1}{32}$
 - (b) $\sqrt{41}$